

TOPOLOGY OF COMPLEX REFLECTION ARRANGEMENTS

DAVID BESSIS

ABSTRACT. Let V be a finite dimensional complex vector space and $W \subset \mathrm{GL}(V)$ be a finite complex reflection group. Let V^{reg} be the complement in V of the reflecting hyperplanes. A classical conjecture predicts that V^{reg} is a $K(\pi, 1)$ space. When W is a complexified real reflection group, the conjecture follows from a theorem of Deligne, [20]. Our main result validates the conjecture for *duality* (or, equivalently, *well-generated*) complex reflection groups. This includes the complexified real case (but our proof is new) and new cases not previously known. We also address a number of questions about $\pi_1(W \setminus V^{\mathrm{reg}})$, the braid group of W .

Let V be a finite dimensional complex vector space and $W \subset \mathrm{GL}(V)$ be an irreducible complex reflection group (all reflection groups considered here are assumed to be finite). Let $d_1 \leq \dots \leq d_n$ be the degrees of W . Let $d_1^* \geq \dots \geq d_n^* = 0$ be the codegrees of W . We say that W is a *duality* group if $d_i + d_i^* = d_n$ for all i (by analogy with the real case, we say that d_n is the *Coxeter number* of W and we often denote it by h). We say that W is *well-generated* if it may be generated by n reflections. Orlik-Solomon observed, by inspecting the classification of Shephard-Todd, that

W is a duality group $\Leftrightarrow W$ is well-generated.

Let V^{reg} be the complement in V of the reflecting hyperplanes. In the case when W is a type A reflection group, Fadell and Neuwirth proved in the early 1960's that V^{reg} is a $K(\pi, 1)$ (this is an elementary use of fibration exact sequences, [22]). Brieskorn conjectured in 1971, [13], that the $K(\pi, 1)$ property holds when W is a complexified real reflection group. It is not clear who first stated the conjecture for arbitrary complex reflection groups. The conjecture may be found in Orlik-Terao's book:

Conjecture 0.1 ([29], p. 163 & p. 259). *The universal cover of V^{reg} is contractible.*

The complexified real case (*i.e.*, Brieskorn's conjecture) was quickly settled by Deligne, [20]. The rank 2 case is trivial. The case of the infinite family was solved in 1983 by Nakamura, [26] (here again, the monomiality of the group allows an efficient use of fibrations). A few other cases immediately follow from the observation by Orlik-Solomon, [28], that certain orbifolds of non-real complex reflection groups are isomorphic to orbifolds of complexified real reflection groups.

Combining all previously known results, the conjecture remained open for six exceptional types: G_{24} , G_{27} , G_{29} , G_{31} , G_{33} and G_{34} . Our main result is:

Theorem 0.2. *Assume that W is a well-generated complex reflection group. The universal cover of V^{reg} is contractible.*

Five of the six open cases are duality groups: G_{24} , G_{27} , G_{29} , G_{33} and G_{34} . The theorem also applies to the complexified real case, for which we obtain a new proof, not relying

on [20]. The only case left is G_{31} , an irreducible complex reflection group of rank 4 which cannot be generated by less than 5 reflections.

Among by-products of our construction, we obtain new cases of several standard conjectures about the braid group of W , defined by

$$B(W) := \pi_1(W \setminus V^{\text{reg}}).$$

Theorem 0.3. *Braid groups of well-generated complex reflection groups are Garside groups.*

In particular, they are torsion-free, have nice solutions to word and conjugacy problems, are biautomatic, admit finite $K(\pi, 1)$ (our construction provides an explicit one), and much more – see [18] for a quite complete reference. None of this was known for the five exceptional groups mentioned above.

Theorem 0.4. *The center of the braid group of an irreducible well-generated complex reflection group is cyclic.*

This, as conjectured by Broué-Malle-Rouquier, [14], should also hold for non-well-generated groups. Again, the cases of G_{24} , G_{27} , G_{29} , G_{33} and G_{34} are new. The only case left is G_{31} .

For $B(G_{29})$, $B(G_{31})$, $B(G_{33})$ and $B(G_{34})$, no presentations were known until now, although some conjectures made in [7] were supported by strong evidences.

Theorem 0.5. *The conjectural presentations for $B(G_{29})$, $B(G_{33})$ and $B(G_{34})$ given in [7] are correct.*

General strategy. The general architecture of our proof is borrowed from Deligne’s original approach but the details are quite different. Every construction here is an analog of a construction from [20] but relies on different combinatorial and geometric objects. Like in [20], one studies a certain braid monoid M , whose structure expresses properties of reduced decompositions in W , and one proves that it is a lattice for the divisibility order (this amounts to saying that the monoid is *Garside*). Like in [20], one uses semi-algebraic geometry to construct an open covering of the universal cover of V^{reg} , with the property that non-empty intersections are contractible. This implies that the universal cover is homotopy equivalent to the nerve of the covering. Like in [20], one interprets this nerve as a certain flag complex obtained from M . Like in [20], the contractibility of the nerve follows from the lattice property for M . However, our proof does not use the classical braid monoid, but a dual braid monoid ([3], [5]), whose construction is generalised to all well-generated complex reflection groups. The construction of the open covering is the most problematic step: by contrast with the real case, one cannot rely on the notions of *walls* and *chambers*. The idea here is to work in $W \setminus V^{\text{reg}}$ and to use a generalisation of the Lyashko-Looijenga morphism. This morphism allows a description of $W \setminus V^{\text{reg}}$ by means of a ramified covering of a type A reflection orbifold. Classical objects like *walls*, *chambers* and *galleries* can somehow be “pulled-back”, via the Lyashko-Looijenga morphism, to give semi-algebraic objects related to the dual braid monoid.

Lyashko-Looijenga coverings. The quotient map $\pi : V^{\text{reg}} \twoheadrightarrow W \setminus V^{\text{reg}}$ is a regular covering. Once a system of basic invariants $(f_1, \dots, f_n)$ is chosen, the quotient space $W \setminus V^{\text{reg}}$

identifies with the complement in $\mathbb{C}^n$ of an algebraic hypersurface $\mathcal{H}$, the *discriminant*, of equation $\Delta \in \mathbb{C}[X_1, \dots, X_n]$. If W is an irreducible duality complex reflection group, it is possible to choose $(f_1, \dots, f_n)$ such that

$$\Delta = X_n^n + \alpha_2 X_n^{n-2} + \dots + \alpha_n,$$

where $\alpha_2, \dots, \alpha_n \in \mathbb{C}[X_1, \dots, X_{n-1}]$. Let $Y := \text{Spec } \mathbb{C}[X_1, \dots, X_{n-1}]$, together with the natural map $p : W \backslash V^{\text{reg}} \rightarrow Y$. We have an identification $W \backslash V \simeq \mathbb{C}^n \xrightarrow{\sim} \mathbb{C} \times Y$ sending the orbit $\bar{v}$ of $v \in V$ to $(f_n(v), p(\bar{v}))$. The fiber of p over $y \in Y$ is a line L_y which intersects $\mathcal{H}$ at n points (counted with multiplicities). Generically, the n points are distinct. Let $\mathcal{K}$ be the *bifurcation locus*, i.e., the algebraic hypersurface of Y consisting of points y such that the intersection has cardinality $< n$. Classical results from invariant theory of complex reflection groups make possible (and very easy) to generalise a construction by Looijenga and Lyashko: the map LL (for “Lyashko-Looijenga”) sending $y \in Y - \mathcal{K}$ to the subset $\{x_1, \dots, x_n\} \subseteq \mathbb{C}$ such that $p^{-1}(y) \cap \mathcal{H} = \{(x_1, y), \dots, (x_n, y)\}$ is a regular covering of degree $n!h^n/|W|$ of the (centered) configuration space of n points in $\mathbb{C}$. In particular, $Y - \mathcal{K}$ is a $K(\pi, 1)$. This observation, which is apparently new in the non-real case, already allows a refinement of our earlier results ([2], [7]) on presentations for the braid group of W .

The dual braid monoid. When W is complexified real, a *dual braid monoid* was constructed in [3] (generalising the construction of Birman-Ko-Lee, [9]; similar partial results were independently obtained by Brady and Brady-Watt, [11]). The construction was later generalised in [5] to the complex reflection group $G(e, e, n)$. Assume W is a well-generated complex reflection group generated by reflections of order 2. Let R be the set of all reflections in W . The idea is that the pair (W, R) has some “Coxeter-like” features. Instead of looking at relations of the type

$$\underbrace{sts\dots}_{m_{s,t}} = \underbrace{tst\dots}_{m_{s,t}},$$

one considers relations of the type

$$st = tu$$

where $s, t, u \in R$. Let $\mathcal{S}$ be the set of all relations of this type holding in W . In general, $B(W) \not\simeq \langle R | \mathcal{S} \rangle$, but it is possible to find natural subsets $R_c \subseteq R$ and $\mathcal{S}_c \subseteq \mathcal{S}$ such that $B(W) \simeq \langle R_c | \mathcal{S}_c \rangle$ (if W is complexified real, $R_c = R$). The elements of $\mathcal{S}_c$ are called *dual braid relations*. The choices of R_c and $\mathcal{S}_c$ are natural once a *Coxeter element* c has been chosen (the notion of *Coxeter element* generalises to the non-real groups in terms of Springer’s theory of regular elements, [34]). Since the relations are positive, one may view the presentation as a monoid presentation, defining a monoid $M(W)$. The crucial property of this monoid is that it is a lattice for the divisibility order or, in other words, a *Garside monoid*. Following Deligne, Bestvina, T. Brady and Charney-Meier-Whittlesey, the Garside structure provides a convenient simplicial Eilenberg-McLane $K(B(W), 1)$ space ([20], [8], [10], [16]). The earlier results on the dual braid monoid are improved here in two directions:

- The construction is generalised to the few exceptional cases $(G_{24}, G_{27}, G_{29}, G_{33}$ and $G_{34})$ not covered by [3] and [5].

- A new geometric interpretation is given, via the Lyashko-Looijenga covering. This interpretation is different for the one given in [3, Section 4].

The second point is the most important. It relies (so far) on a counting argument, following and extending a property which, for the complexified real case, was conjectured by Looijenga and proved in a letter from Deligne to Looijenga, [21].

Tunnels. The classical theory of real reflection groups combines a “combinatorial” theory (Coxeter systems) and a “geometric” theory (expressed in the language, invented by Tits, of walls, chambers, galleries, buildings...) We expect the dual braid monoid approach to eventually provide effective substitutes for much of this classical theory. We use here a notion of *tunnel*, which is a rudimentary geometric object replacing the classical notion of *minimal gallery between two chambers*. An important difference with the classical geometric language is that tunnels are naturally visualised in $W \backslash V$ (instead of V). A tunnel T is a path in $W \backslash V^{\text{reg}}$ drawn inside a single line L_y (for some $y \in Y$) and with constant imaginary part. It represents an element b_T of the dual braid monoid M . An element of M is *simple* if it is represented by a tunnel. This notion coincides with the notion of simple element associated with the Garside structure. In the classical approach, for any chamber $\mathcal{C}$, there are as many equivalence classes of minimal galleries starting at $\mathcal{C}$ as simple elements (this number is $|W|$). Here the situation is different: in a given L_y , not all simples are represented. The simples represented in different L_y ’s may be compared thanks to a huge “basepoint” $\mathcal{U}$ which is both dense in $W \backslash V^{\text{reg}}$ and contractible.

Non-crossing partitions. In the classical cases A_n , B_n , D_n and, more generally, $G(e, e, n)$, the structure of $M(W)$ is understood in terms of suitable notions of *non-crossing partitions* ([1], [5], [6]). The dual braid monoid of an irreducible well-generated complex reflection W gives rise to a lattice of *generalised non-crossing partitions*, whose cardinal is the *generalised Catalan number*

$$\text{Cat}(W) := \prod_{i=1}^n \frac{d_i + d_n}{d_i}.$$

(the term “partition” should not be taken too seriously: the elements of the lattice no longer have a natural interpretation as partitions.) It is likely that this combinatorial object has some representation-theoretic interpretation. If W is not well-generated, $\text{Cat}(W)$ may fail to be an integer.

Acknowledgments. Work on this project started during stimulating visits to KIAS (Seoul) and RIMS (Kyoto), in the summer of 2003, when I realised that the dual braid monoid construction extends to duality groups. I thank Sang Jin Lee and Kyoji Saito for their hospitality and for their interests in discussing these subjects. After this initial progress, I remained stucked for many months, trying to construct the open covering using affine geometry in V^{reg} (refining [3, Section 4]). Two observations helped me decide to work in the quotient. First, Kyoji Saito pointed out that the starting point for the construction of the *flat structure* (or *Frobenius manifold structure*, see [32]) on real reflection orbifolds was precisely the duality between degrees and codegrees. The intuition that the flat structure has something to do with the $K(\pi, 1)$ property is explicitly mentioned as a motivation for [32] (see also [33, p. 7]). The second useful discussion was with Frédéric

Chapoton, who pointed out the numerological coincidence, in the Coxeter case, between the degree of the Lyashko-Looijenga covering and the number of maximal chains in the lattice of non-crossing partitions.

Warning. This text was initially planned to be a research announcement. Though it grew much beyond any reasonable size for a research announcement, it is not yet in final form. It still misses some details, and several arguments are not optimal.

NOTATIONS

We save the letter i for indexing purposes and denote by $\sqrt{-1}$ a complex square root of -1 fixed once for all. If n is a positive integer, we denote by ζ_n the standard n -th root of unity $\exp(2\sqrt{-1}\pi/n)$.

Many objects depend on a complex reflection group W , e.g., the braid group $B(W)$. We often drop the explicit mention of W , and write B for $B(W)$. When n is an integer, we denote by B_n the braid group on n strings, together with its standard generating set $\sigma_1, \dots, \sigma_{n-1}$; it is isomorphic to the braid group of $\mathfrak{S}_n$ in its permutation reflection representation (see Section 3). The groups B and B_n appear simultaneously and should not be confused.

TOPOLOGICAL CONVENTIONS

Let E be a topological space. Let γ be a path in E , *i.e.*, a continuous map $[0, 1] \rightarrow E$. We say that γ is a *path from $\gamma(0)$ to $\gamma(1)$* – or that $\gamma(0)$ is the *source* and $\gamma(1)$ the *target*. The concatenation rule is as follows: if γ, γ' are paths such that $\gamma(1) = \gamma'(0)$, we define by $\gamma \circ \gamma'$ the path mapping $t \leq 1/2$ to $\gamma(2t)$ and $t \geq 1/2$ to $\gamma'(2t - 1)$.

A path γ from e to e' induces an isomorphism $\phi_\gamma : \pi_1(E, e) \xrightarrow{\sim} \pi_1(E, e')$.

Let X be a subspace of E . Assume that X is simply-connected (in all applications, X will actually be contractible). Then for all $x, x' \in X$ there is a unique homotopy class of paths from x to x' drawn in X , thus a natural isomorphism $\phi_{X,x,x'} : \pi_1(E, x) \xrightarrow{\sim} \pi_1(E, x')$. By unicity, $\phi_X := (\phi_{X,x,x'})_{x,x' \in X}$ is a transitive system of isomorphisms between the $(\pi_1(E, x))_{x \in X}$. We denote by

$$\pi_1(E, X)$$

the transitive limit. Practically speaking, $\pi_1(E, X)$ should be thought of as any $\pi_1(E, x)$, for some $x \in X$, together with an unambiguous recipe thanks to which any path in E from any $x' \in X$ to any $x'' \in X$ represents a unique element of $\pi_1(E, x)$ – moreover, one may forget about which $x \in X$ was chosen and change it at our convenience.

Assume that E is locally simply-connected. To get a model $\tilde{E}_x$ of the universal cover $\tilde{E}$, one chooses a basepoint $x \in E$. A path γ from x to $e \in E$ represents a point $\bar{\gamma} \in \tilde{E}_x$ in the fiber over e . Two paths represent the same point if and only if they are homotopic (with fixed endpoints). The topology of $\tilde{E}_x$ around $\bar{\gamma}$ is described by concatenation at the target with paths starting at e and drawn in simply-connected neighbourhoods of e . The fundamental group $\pi_1(E, x)$ acts on the left on $\tilde{E}_x$ by concatenation at the source. More generally, a path from x to x' induces an isomorphism $\tilde{E}_x \xrightarrow{\sim} \tilde{E}_{x'}$. Here again, if X is simply-connected, we have a transitive system of isomorphisms between $(\tilde{E}_x)_{x \in X}$ whose limit is a model $\tilde{E}_X$ of the universal cover. Practically speaking, any path from any $x \in X$

to any $e \in E$ unambiguously represents a point in $\tilde{E}_X$ (in the fiber over e). Moreover, $\pi_1(E, X)$ acts on the left on $\tilde{E}_X$.

1. COMPLEX REFLECTION GROUPS, DISCRIMINANTS, BRAID GROUPS

Let V be a $\mathbb{C}$ -vector space of finite dimension n . A *complex reflection group* in $\mathrm{GL}(V)$ is a (finite) subgroup generated by *complex reflection*, elements whose fixed subspace is a hyperplane. Let $W \subset \mathrm{GL}(V)$ be a complex reflection group. A *system of basic invariants* is an n -tuple $f = (f_1, \dots, f_n)$ of homogeneous W -invariant polynomial functions on V generating the algebra $\mathbb{C}[V]^W$ of W -invariant polynomial functions. A classical theorem of Shephard-Todd, [35], asserts that such tuples exist, and that they consist of algebraically independent terms. Set $d_i := \deg f_i$; these numbers are the *degrees* of W . Up to reordering, we may assume that $d_1 \leq d_2 \leq \dots \leq d_n$. The sequence $(d_1, \dots, d_n)$ is then independent of the choice of f .

Choosing a system of basic invariants f amounts to choosing a graded algebra isomorphism $\mathbb{C}[V]^W \simeq \mathbb{C}[X_1, \dots, X_n]$, $f_i \mapsto X_i$, where the indeterminate X_i is declared homogeneous with degree d_i . Geometrically, this isomorphism identifies the categorical quotient $W \backslash V$ with the affine space $\mathbb{C}^n$.

Further features of the invariant theory of complex reflection groups involve invariant vector fields and invariant differential forms on V .

Theorem 1.1 ([29], Lemma 6.48). *The $\mathbb{C}[V]^W$ -modules $(\mathbb{C}[V] \otimes V)^W$ and $(\mathbb{C}[V] \otimes V^*)^W$ are free of rank n .*

If $f = (f_1, \dots, f_n)$ is a system of basic invariants, $df := (df_1, \dots, df_n)$ is a $\mathbb{C}[V]^W$ -basis for $(\mathbb{C}[V] \otimes V^*)^W$. Being homogeneous, the module $(\mathbb{C}[V] \otimes V)^W$ admits an homogeneous basis.

Definition 1.2. A system of basic derivations is an homogeneous $\mathbb{C}[V]^W$ -basis $\xi = (\xi_1, \dots, \xi_n)$ of $(\mathbb{C}[V] \otimes V)^W$, with $\deg(\xi_1) \geq \deg(\xi_2) \geq \dots \geq \deg(\xi_n)$.

The sequence $(d_1^*, \dots, d_n^*) := (\deg(\xi_1), \dots, \deg(\xi_n))$ (which does not depend on the choice of ξ) is the sequence of *codegrees* of W .

Note that, as in [2], we label codegrees in decreasing order, which is slightly unusual. When W is a complexified real reflection group, we have $V \simeq V^*$ as W -modules, thus $d_i^* = d_{n-i+1} - 2$ for all i . This relation is specific to the real situation and is not relevant here.

The Euler vector field on V is invariant and of degree 0. Thus $d_n^* = 0$.

Invariant vector fields define vector fields on the quotient variety. Let f be a system of basic invariants and ξ be a system of basic derivations. For $j \in \{1, \dots, n\}$, the vector field ξ_j defines a vector field $\bar{\xi}_j$ on $W \backslash V$. Since $\frac{\partial}{\partial f_1}, \dots, \frac{\partial}{\partial f_n}$ is a $\mathbb{C}[V]^W$ -basis of the module of polynomial vector fields on $W \backslash V$, we have

$$\bar{\xi}_j = \sum_{i=1}^n m_{i,j} \frac{\partial}{\partial f_i},$$

where the $m_{i,j}$ are uniquely defined elements of $\mathbb{C}[V]^W$.

Definition 1.3. The discriminant matrix of W (with respect to f and ξ) is $M := (m_{i,j})_{i,j}$.

To recover ξ_j from $\overline{\xi_j}$, replace $\frac{\partial}{\partial f_i}$ by $\sum_{k=1}^n \frac{\partial f_i}{\partial Y_k} \frac{\partial}{\partial Y_i}$. By weighted homogeneity, one has:

Lemma 1.4. *For all i, j , $\text{wt}(m_{i,j}) = d_i + d_j^*$.*

The vector space V decomposes as a direct sum $\bigoplus_i V_i$ of irreducible representations of W . Denote by W_i the irreducible reflection group in $\text{GL}(V_i)$ generated by (the restriction of) the reflections in W whose hyperplanes contain $\bigoplus_{j \neq i} V_j$. We have $W \simeq \prod_i W_i$. Viewing W and the W_i 's as *reflection groups*, i.e., groups endowed with a reflection representation, it is natural to actually write $W = \bigoplus_i W_i$.

We denote by $\mathcal{A}$ the arrangement of W , i.e., the set of reflecting hyperplanes of reflections in W ; it is in bijection with the disjoint union $\sqcup_i \mathcal{A}_i$, where $\mathcal{A}_i$ is the arrangement of W_i . We set

$$V^{\text{reg}} := V - \bigcup_{H \in \mathcal{A}} H.$$

Denote by p the quotient map $V \rightarrow W \backslash V$. Choose a basepoint $v_0 \in V^{\text{reg}}$.

Definition 1.5 ([14]). The *braid group* of W is $B(W) := \pi_1(W \backslash V^{\text{reg}}, p(v_0))$.

To write explicit equations, one chooses a system of basic invariants f . The *discriminant* $\Delta(W, f) \in \mathbb{C}[X_1, \dots, X_n]$ is the reduced equation of $p(\bigcup_{H \in \mathcal{A}} H)$, via the identification $\mathbb{C}[V]^W \simeq \mathbb{C}[X_1, \dots, X_n]$.

One easily sees that $B(W) \simeq \prod_i B(W_i)$. More generally, all objects studied here behave “semi-simply”, and we may restrict our attention to irreducible complex reflection groups.

Since $B(W)$ is the fundamental group of the complement of an algebraic hypersurface, it is generated by particular elements called *generators-of-the-monodromy* or *meridiens* (see, for example, [14] or [2]). They map to reflections under the natural epimorphism $B(W) \rightarrow W$. The diagrams given in [14] symbolise presentations whose generators are generators-of-the-monodromy (except for the six exceptional types for which no presentation was known).

Definition 1.6. The generators-of-the-monodromy of $B(W)$ are called *braid reflections*.

This terminology was suggested by Broué. It is actually tempting to simply call them *reflections*: since they generate $B(W)$, the braid group appears to be some sort of “reflection group”.

Another natural feature of $B(W)$ is the existence of a natural length function, which is the unique group morphism

$$l : B(W) \rightarrow \mathbb{Z}$$

such that, for all braid reflection $s \in B(W)$, $l(s) = 1$.

Consider the intersection lattice $\mathcal{L}(\mathcal{A}) := \{\bigcap_{H \in \mathcal{A}} H \mid A \subseteq \mathcal{A}\}$. Elements of $\mathcal{L}(\mathcal{A})$ are called *flats*. It is standard to endow $\mathcal{L}(\mathcal{A})$ with the reversed-inclusion partial ordering:

$$\forall L, L' \in \mathcal{L}(\mathcal{A}), L \leq L' :\Leftrightarrow L \supseteq L'.$$

For $L \in \mathcal{L}(\mathcal{A})$, we denote by L^0 the complement in L of the flats strictly included in L . The $(L^0)_{L \in \mathcal{L}(\mathcal{A})}$ form a stratification $\mathcal{S}$ of V . We consider the partial ordering on $\mathcal{S}$ defined by $L^0 \leq L'^0 :\Leftrightarrow L \leq L'$. This is a degeneracy relation:

$$\forall L \in \mathcal{L}(\mathcal{A}), \overline{L^0} = L = \bigcup_{S \in \mathcal{S}, L^0 \leq S} S.$$

Since W acts on $\mathcal{A}$, it acts on $\mathcal{L}(\mathcal{A})$ and we obtain a quotient stratification $\overline{\mathcal{S}}$ of $W \backslash V$ called *discriminant stratification*.

Proposition 1.7 ([29], Corollary 6.114). *Let $v \in V$. The vectors $\xi_1(v), \dots, \xi_n(v)$ span the tangent space to the stratum of $\mathcal{S}$ containing v . The vectors $\bar{\xi}_1(v), \dots, \bar{\xi}_n(v)$ span the tangent space to the stratum of $\overline{\mathcal{S}}$ containing $\bar{v}$.*

2. WELL-GENERATED COMPLEX 2-REFLECTION GROUPS

Irreducible complex reflection groups were classified fifty years ago by Shephard and Todd, [35]. There is an infinite family $G(de, e, n)$, where d, e, n are positive integers, and 34 exceptions $G_4, \dots, G_{37}$. Let us distinguish three subclasses of complex reflection groups:

- *real reflection groups*, obtained by scalar extension from reflection groups of real vector spaces;
- *2-reflection groups*, generated by reflections of order 2;
- *well-generated reflection groups*, i.e., irreducible complex reflection groups generated by n reflections, where n is the dimension, and more generally direct sums of such irreducible groups.

Real reflection groups are both 2-reflection groups and well-generated. For non-real groups, any combination of the other two properties may hold.

As far as the $K(\pi, 1)$ conjecture and properties of braid groups are concerned, it is enough to restrict one's attention to 2-reflection groups, thanks to the following theorem.

Theorem 2.1. *For any complex reflection group $W \subset \mathrm{GL}(V)$, one may find V' a complex vector space, $W' \subset \mathrm{GL}(V')$ a complex 2-reflection group and f (resp. f') a system of basic invariants for W (resp. W') such that $\Delta(W, f) = \Delta(W', f')$. In particular, $W \backslash V^{\mathrm{reg}} \simeq W' \backslash V'^{\mathrm{reg}}$ and $B(W) \simeq B(W')$.*

Proof. This theorem may be observed on the classification and was certainly known to experts but remained unexplained. The recent work of Couwenberg-Heckman-Looijenga, [17], may be adapted to provide a direct argument, which is sketched below.

All references and notations are from *loc. cit.*. Assume that W is not a 2-reflection group. For each $H \in \mathcal{A}$, let e_H be the order of the pointwise stabiliser W_H and set $\kappa_H := 1 - e_H/2$. Consider the Dunkl connection ∇ with connection form $\sum_{H \in \mathcal{A}} \omega_H \otimes \kappa_H \pi_H$, as in Example 2.5. Since $e_H \geq 2$, we have $\kappa_H \leq 0$. In particular, $\kappa_0 = 1/n \sum_{H \in \mathcal{A}} \kappa_H \leq 0$ (Lemma 2.13) and we are in the situation of *loc. cit.* Section 5. In many cases, this suffices to conclude. The problem is that, even though at least some e_H have to be > 2 , it is possible that $\mathcal{A}$ contains several orbits, some of them with $e_H = 2$. To handle this, one has to enlarge the ‘‘Schwarz symmetry group’’ of *loc. cit.*, Section 4. $\square$

The importance of the distinction between well-generated and ‘‘badly-generated’’ groups was first pointed out by Orlik-Solomon, who observed in [27] a coincidence with invariant-theoretical aspects.

Theorem 2.2. *Let W be an irreducible complex reflection group. The following assertions are equivalent:*

- (i) W is well-generated.

- (ii) For all $i \in \{1, \dots, n\}$, $d_i + d_i^* = d_n$.
- (iii) For all $i \in \{1, \dots, n\}$, $d_i + d_i^* \leq d_n$.
- (iv) For any system of basic invariants f , there exists a system of basic derivations ξ such that the discriminant matrix decomposes as $M = M_0 + X_n M_1$, where M_0, M_1 are matrices with coefficients in $\mathbb{C}[X_1, \dots, X_{n-1}]$ and M_1 is lower triangular with non-zero scalars on the diagonal.
- (v) For any system of basic invariants f , we have $\frac{\partial^n \Delta(W, f)}{(\partial X_n)^n} \in \mathbb{C}^\times$ (in other words, $\Delta(W, f)$, viewed as a polynomial in X_n with coefficients in $\mathbb{C}[X_1, \dots, X_{n-1}]$, is monic of degree n).

The matrix M_1 from assertion (iv) is an analog of the matrix J^* from [32], p. 10. Assertion (iv) itself generalises the non-degeneracy argument for J^* , which is an important piece of the construction of Saito's "flat structure".

Proof. (i) $\Rightarrow$ (ii) was observed in [27] inspecting the classification. We still have no good explanation.

(ii) $\Rightarrow$ (iii) is trivial.

(iii) $\Rightarrow$ (v). Let $h := d_n$. A first step is to observe that, under assumption (iii), h is a regular number, in the sense of Springer, [34]. Indeed, since d_n^* is the only codegree equal to 0, the condition $d_i + d_i^* \leq h$ implies, for $i = 1, \dots, n-1$, both $0 < d_i < h$ and $0 < d_i^* < h$; in particular, except when $i = n$, h does not divide neither d_i nor d_i^* . A criterion first noticed by Lehrer-Springer and proved abstractly by Lehrer-Michel ([24]) says that if a number d divides exactly as many degrees as codegrees, then it is regular. Since h is regular and divides only one degree, we may use [2], Lemma 1.6 (ii) to obtain assertion (v): the discriminant is X_n -monic, and by weighted-homogeneity it must be of degree n .

(iii) $\Rightarrow$ (iv) is a refinement of the previous discussion. Each entry $m_{i,j}$ of the matrix M is weighted-homogeneous of total weight $d_i + d_j^* \leq d_n + d_1^* = 2h - d_1 < 2h$; since X_n has weight h , $\deg_{X_n} m_{i,j} \leq 1$. This explains the decomposition $M = M_0 + X_n M_1$, where M_0 and M_1 have coefficients in $\mathbb{C}[Y]$. If $i < j$ and $d_i < d_j$, then $d_i + d_j^* < d_j + d_j^* = h$, thus $\deg_{X_n} m_{i,j} = 0$. The matrix M_0 is *weakly lower triangular*, i.e., lower triangular except that there could be non-zero terms above the diagonal in square diagonal blocks corresponding to successive equal degrees (successive degrees may indeed be equal, as in the example of type D_4 , where the degrees are 2, 4, 4, 6).

Let $i_0 < j_0$ such that $d_{i_0} = d_{i_0+1} = \dots = d_{j_0}$ (looking at the classification, one may observe that this forces $j = i + 1$; this observation is not used in the argument below). For all $i, j \in \{i_0, \dots, j_0\}$, we have $d_i + d_j^* = h$. By weighted homogeneity, this implies that the corresponding square block of M_1 consists of scalars. The basic derivations $\xi_{i_0}, \dots, \xi_{j_0}$ all have the same degrees, thus one is allowed to perform Gaussian elimination on the corresponding columns of M . Thus, up to replacing ξ by another system of basic derivations ξ' , we may assume that M_1 is lower triangular.

The diagonal terms of M_1 must be scalars, once again by weighted homogeneity. Assuming (iii), we already know that (v) holds. The determinant of M is $\Delta(X, f)$; (v) implies that the coefficient of X_n^n is non-zero. This coefficient is the product of the diagonal terms of M_1 . We have proved (iii).

(iv) $\Rightarrow$ (v) is trivial.

(v) $\Rightarrow$ (i) follows from the main result in [2]. □

The following notion was considered in [31] for real reflection groups.

Definition 2.3. A system of basic derivations is *flat* (with respect to f) if the discriminant matrix may be written $M = M_0 + X_n \text{Id}$, where M_0 is a matrix with coefficients in $\mathbb{C}[X_1, \dots, X_{n-1}]$.

Theorem 2.2 has the following corollary:

Corollary 2.4. *Let W be an well-generated irreducible reflection group. The flatness of a system of basic derivations does not depend from the choice of f . There exists flat systems of basic derivations.*

Proof. The independence follows from the fact that $d_{n-1} < d_n$, which itself follows from characterisation (ii) of well-generated groups.

Existence: choose f and ξ , and write $M = M_0 + X_n M_1$, as in characterisation (iv). The matrix M_1 is invertible in $\text{GL}_n(\mathbb{C}[X_1, \dots, X_{n-1}])$. The matrix $M_1^{-1}M = M_1^{-1}M_0 + X_n \text{Id}$ represents a flat system of basic derivations (weighted homogeneity is preserved by the Gaussian elimination procedure). $\square$

Contrary to what happens with real reflection groups, we may not use the identification $V \simeq V^*$ to obtain “flat system of basic invariants”.

Irreducible groups which are not well-generated may always be generated by $\dim V + 1$ reflections. This fact has been observed long ago, by case-by-case inspection, but no general argument is known. (In some sense, these badly-generated groups should be thought of as affine groups). The simplest example of a non-well-generated group is the group $G(4, 2, 2)$, generated by

$$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}.$$

In the sequel, several arguments are case-by-case. As explained above, one has to consider only well-generated irreducible complex 2-reflection groups. Their list consists of:

- the Coxeter types $A_n, B_n, D_n, E_6, E_7, E_8, F_4, H_3, H_4$ and $I_2(e)$,
- the infinite family $G(e, e, n)$, where e and n are integers ≥ 2 , which interpolates the Coxeter series D_n (when $e = 2$) and $I_2(e)$ (when $n = 2$),
- five non-real exceptional types $G_{24}, G_{27}, G_{29}, G_{33}$ and G_{34} .

3. SYMMETRIC GROUPS, CONFIGURATIONS SPACES AND CLASSICAL BRAID GROUPS

This section is only included to clarify some terminology and notations. Everything here is classical and elementary.

Let n be a positive integer. The symmetric group $\mathfrak{S}_n$ may be viewed as reflection group, acting on $\mathbb{C}^n$ by permuting the canonical basis. This representation is not irreducible. Let H be the hyperplane of equation $\sum_{i=1}^n X_i$ (where $X_1, \dots, X_n$ is the dual canonical basis of $\mathbb{C}^n$). It is preserved by $\mathfrak{S}_n$, which acts on it as an irreducible complex reflection group.

We have $\mathbb{C}[X_1, \dots, X_n]^{\mathfrak{S}_n} = \mathbb{C}[\sigma_1, \dots, \sigma_n]$ and $\mathbb{C}[H]^{\mathfrak{S}_n} = \mathbb{C}[\sigma_1, \dots, \sigma_n]/\sigma_1$, where $\sigma_1, \dots, \sigma_n$ are the elementary symmetric functions on $X_1, \dots, X_n$. Set $E'_n := \mathfrak{S}_n \backslash \mathbb{C}^n = \text{Spec } \mathbb{C}[\sigma_1, \dots, \sigma_n]$

and $E_n := \mathfrak{S}_n \backslash H = \text{Spec } \mathbb{C}[\sigma_1, \dots, \sigma_n] / \sigma_1$. These spaces have more convenient descriptions in terms of multisets.

Recall that a *multiset* is a set S (the *support* of the multiset) together with a map $m : S \rightarrow \mathbb{Z}_{\geq 1}$ (the *multiplicity*). The *cardinal* of such a multiset is $\sum_{s \in S} m(s)$ (it lies in $\mathbb{Z}_{\geq 0} \cup \{\infty\}$). If (S, m) and (S', m') are two multisets and if S, S' are subsets of a common ambient set, then we may define a *multiset (disjoint) union* $(S, m) \cup (S', m')$, whose support is $S \cup S'$ and whose multiplicity is $m + m'$ (where m , resp. m' , is extended by 0 outside S , resp. S').

Let $(x_1, \dots, x_n) \in \mathbb{C}^n$. The associated $\mathfrak{S}_n$ -orbit is uniquely determined by the multiset $\{\{x_1, \dots, x_n\}\} := \bigcup_{i=1}^n (\{x_i\}, 1)$. This identifies E'_n with the set of multisets of cardinal n with support in $\mathbb{C}$ (such multisets are called *configurations of n points in $\mathbb{C}$*). The subvariety E_n , defined by $\sigma_1 = 0$, consists of configurations $\{\{x_1, \dots, x_n\}\}$ with $\sum_{i=1}^n x_i = 0$ (such configurations are *centered*). The natural inclusion $E_n \subseteq E'_n$ admits the retraction ρ defined by

$$\rho(\{\{x_1, \dots, x_n\}\}) := \left\{ \left\{ x_1 - \sum_{i=1}^n x_i / n, \dots, x_n - \sum_{i=1}^n x_i / n \right\} \right\}.$$

Algebraically, this corresponds to the identification of $\mathbb{C}[\sigma_1, \dots, \sigma_n] / \sigma_1$ with $\mathbb{C}[\sigma_2, \dots, \sigma_n]$.

We find it convenient to use configurations in E'_n to represent elements of E_n , implicitly working through ρ . E.g., in the proof of Proposition 8.3, it makes sense to describe a deformation retraction of a subspace of E_n to a point in terms of arbitrary configurations because the construction, which only implies the relative values of the x_i 's, is compatible with ρ . We adopt this viewpoint from now on, without further justifications (compatibility will always be obvious).

Consider the lexicographic total ordering of $\mathbb{C}$: if $z, z' \in \mathbb{C}$, we set

$$z \leq z' :\Leftrightarrow \begin{cases} \text{re}(z) < \text{re}(z') & \text{or} \\ \text{re}(z) = \text{re}(z') \text{ and } \text{im}(z) \leq \text{im}(z'). \end{cases}$$

Definition 3.1. The *ordered support* of an element of E_n is the unique sequence $(x_1, \dots, x_k)$ such that $\{x_1, \dots, x_k\}$ is the support and $x_1 < x_2 < \dots < x_k$.

We may uniquely represent an element of E_n by its ordered support $(x_1, \dots, x_k)$ and the sequence $(n_1, \dots, n_k)$ of multiplicities at $x_1, \dots, x_k$.

The regular orbit space $E_n^{\text{reg}} := \mathfrak{S}_n \backslash H^{\text{reg}}$ consists of those multisets whose support has cardinal n (or, equivalently, whose multiplicity is constantly equal to 1). More generally, the strata of the discriminant stratification of E_n are indexed by partitions of n : the stratum S_λ associated with a partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$, where the λ_j 's are integers with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k > 0$ and $\sum_{j=1}^k \lambda_j = n$, consists of configurations whose supports has cardinal k and whose multiplicity functions take the values $\lambda_1, \dots, \lambda_k$ (with multiplicities).

The braid group B_n associated with $\mathfrak{S}_n$ is the usual braid group on n strings. We need to be more precise about our choice of basepoint. For this purpose, we define

$$E_n^{\text{gen}}$$


as the subset of E_n^{reg} consisting of configurations of n points with distinct real parts (this is the first in a series of definitions of semi-algebraic nature). It is clear that:

Lemma 3.2. E_n^{gen} is contractible.

Using our topological conventions, we set

$$B_n := \pi_1(E_n, E_n^{\text{gen}}).$$

This group admits a *standard generating set* (the one considered by Artin), consisting of braid reflections $\sigma_1, \dots, \sigma_{n-1}$ defined as follows (we used bold fonts to avoid confusion with the elementary symmetric functions). Let $(x_1, \dots, x_n)$ be the ordered support of a point in E_n^{gen} . Then σ_i is represented by the following motion of the support:



Artin's presentation for B_n is

$$B_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \sigma_i \sigma_j = \sigma_j \sigma_i \text{ if } |i - j| > 1 \rangle.$$

The following definition requires a compatibility condition which is a classical elementary consequence of the above presentation.

Definition 3.3. Let G be a group. The (*right*) *Hurwitz action* of B_n on G^n is defined by

$$(g_1, \dots, g_{i-1}, g_i, g_{i+1}, g_{i+2}, \dots, g_n) \cdot \sigma_i := (g_1, \dots, g_{i-1}, g_{i+1}, g_{i+1}^{-1} g_i g_{i+1}, g_{i+2}, \dots, g_n),$$

for all $(g_1, \dots, g_n) \in G^n$ and all $i \in \{1, \dots, n-1\}$.

This action preserves the fibers of the product map $G^n \rightarrow G, (g_1, \dots, g_n) \mapsto g_1 \dots g_n$.

4. LYASHKO-LOOIJENGA COVERINGS

Let W be an irreducible well-generated complex reflection group, together with a system of basic invariants f and a flat system of basic derivations ξ with discriminant matrix $M_0 + X_n \text{Id}$. Expanding the determinant, we observe that

$$\Delta_f = \det(M_0 + X_n \text{Id}) = X_n^n + \alpha_2 X_n^{n-2} + \alpha_3 X_n^{n-3} + \dots + \alpha_n,$$

where $\alpha_i \in \mathbb{C}[Y]$. Since Δ_f is weighted homogeneous of total weight nh for the system of weights $\text{wt}(X_i) = d_i$, each α_i is weighted homogeneous of weight ih .

Definition 4.1. The (generalised) *Lyashko-Looijenga morphism* is the morphism LL from $Y = \text{Spec } \mathbb{C}[X_1, \dots, X_{n-1}]$ to $E_n \simeq \text{Spec } \mathbb{C}[\sigma_2, \dots, \sigma_n]$ defined by $\sigma_i \mapsto (-1)^i \alpha_i$.

Of course, we have in mind the geometric interpretation of LL explained below. Let $v \in V$. The orbit $\bar{v} \in W \backslash V$ is represented by a pair

$$(x, y) \in \mathbb{C} \times Y,$$

where $x = f_n(v)$ and y is the point in Y with coordinates $(f_1(v), \dots, f_{n-1}(v))$ (this coding of points in $W \backslash V$ will be used throughout this article). Denote by p the projection $W \backslash V \rightarrow Y, (x, y) \mapsto y$.

Definition 4.2. For any point y in Y , we denote by L_y the fiber of the projection $p : W \backslash V \rightarrow Y$ over y .

For any $y \in Y$, the affine line L_y intersects the discriminant $\mathcal{H}$ in n points (counted with multiplicities), whose coordinates are

$$(x_1, y), \dots, (x_n, y),$$

where $\{\{x_1, \dots, x_n\}\}$ is the multiset of solutions in X_n of $\Delta_f = 0$ where each α_i has been replaced by its value at y . We have

$$LL(y) = \{\{x_1, \dots, x_n\}\}.$$

Definition 4.3. The *bifurcation locus* is the algebraic hypersurface $\mathcal{K} \subseteq Y$ defined by the equation $\text{Disc}_{X_n}(\Delta_f) = 0$.

By $\text{Disc}_{X_n}(\Delta_f)$, we mean the discriminant of Δ_f , seen as a one-variable polynomial with coefficients in $\mathbb{C}[Y]$.

The first theorem of this section is an (elementary) extension to our context of a result from [25].

Theorem 4.4. *The morphism LL is a ramified covering of degree $n!h^n/|W|$. It restricts to an unramified covering $Y - \mathcal{K} \rightarrow E_n^{\text{reg}}$.*

The first lemma generalises [25, Theorem 1.4]:

Lemma 4.5. *LL is étale on $Y - \mathcal{K}$.*

Proof. As mentioned in [25, (1.5)], the lemma will follow if we prove that for all $y \in Y - \mathcal{K}$ with $LL(y) = \{x_1, \dots, x_n\}$, the hyperplanes $H_1, \dots, H_n$ tangent to $\mathcal{H}$ at the n distinct points $(x_1, y), \dots, (x_n, y)$ are in general position.

To prove this, we use Proposition 1.7. Each H_i is spanned by $\bar{\xi}_1(x_i, y), \dots, \bar{\xi}_n(x_i, y)$. Let $(\varepsilon_1, \dots, \varepsilon_n)$ be the basis of $(W \setminus V)^*$ dual to $(\frac{\partial}{\partial X_1}, \dots, \frac{\partial}{\partial X_n})$. Let $l_i = \sum_j \lambda_{j,i} \varepsilon_j$ be a non-zero vector in $(W \setminus V)^*$ orthogonal to H_i . This amounts to taking a non-zero column vector $(\lambda_{j,i})_{j=1, \dots, n}$ in the kernel of $M(x_i, y)$, or equivalently an eigenvector of $M_0(y)$ associated to the eigenvalue $-x_i$. By assumption, the x_i are distinct. The eigenvectors are linearly independent. $\square$

The theorem then follows from:

Lemma 4.6. *LL is finite, of degree $n!h^n/|W|$.*

Proof. Using the first lemma, we observe that $\alpha_2, \dots, \alpha_n$ are algebraically independent. The lemma follows from a standard argument from the theory of quasi-homogeneous morphisms. Since each X_i has weight d_i , the weighted Poincaré series of $\mathbb{C}[Y] = \mathbb{C}[X_1, \dots, X_{n-1}]$ is $\prod_{i=1}^{n-1} \frac{1}{1-t^{d_i}}$. The weighted Poincaré series of $\mathbb{C}[\alpha_2, \dots, \alpha_n]$ is $\prod_{i=2}^n \frac{1}{1-t^{ih}}$. The extension is finite, of degree

$$\prod_{i=1}^{n-1} \frac{(i+1)h}{d_i} = \frac{n!h^{n-1}}{d_1 \dots d_{n-1}} = \frac{n!h^n}{|W|}.$$

$\square$

Corollary 4.7. *The space $Y - \mathcal{K}$ is a $K(\pi, 1)$.*

This corollary suffices to obtain one of the results mentionned in the introduction, Theorem 0.5. Indeed, it was observed in the last paragraph of [7, section 4] that proving that $\pi_2(Y - \mathcal{K}) = 0$ was sufficient to turn our conjectures into theorems. At this stage, the theorem still relies on the brutal computations described in *loc. cit.*. A more direct approach is possible:

Since $\pi_2(Y - \mathcal{K}) = 0$, we may directly compute a presentation for B using a Van Kampen type method. The map $p : W \setminus\!\! \setminus V^{\text{reg}} \rightarrow Y$ restricts to a fibration, still denoted by p , between $E := p^{-1}(Y - \mathcal{K})$ and $Y - \mathcal{K}$. Choose a basepoint $y \in Y - \mathcal{K}$, let F be the fiber $L_y \cap W \setminus\!\! \setminus V$, choose a basepoint $(x, y) \in F$. Consider the following commutative diagram, whose first line is the end of the fibration exact sequence, and where β comes from the inclusion of spaces:

$$\begin{array}{ccccccc} \pi_2(Y - \mathcal{K}, y) = 1 & \longrightarrow & \pi_1(F, (x, y)) & \xrightarrow{\iota} & \pi_1(E, (x, y)) & \xrightarrow{p_*} & \pi_1(Y - \mathcal{K}, y) \longrightarrow 1 \\ & & & \searrow \alpha & \downarrow \beta & & \\ & & & & \pi_1(W \setminus\!\! \setminus V^{\text{reg}}, (x, y)). & & \end{array}$$

We are in the context of [2, Theorem 2.5], from which we conclude that α is surjective. The space E is the complement in $W \setminus\!\! \setminus V \simeq \mathbb{C}^n$ of the hypersurface $\mathcal{H} \cup p^{-1}(\mathcal{K})$, where $\mathcal{H}$ is the discriminant (defined by $\Delta(W, f) = 0$). The space $W \setminus\!\! \setminus V^{\text{reg}}$ is the complement of $\mathcal{H}$. By [2, Lemma 2.1], the kernel of β is generated by the generators-of-the-monodromy around $p^{-1}(\mathcal{K})$.

Let us prove that the exact sequence in the first line of the diagram is split. Let $\gamma_1, \dots, \gamma_m$ be generators-of-the-monodromy in $\pi_1(Y - \mathcal{K}, y)$ such that $\{\gamma_1, \dots, \gamma_m\}$ generates $\pi_1(Y - \mathcal{K}, y)$. Let $\mathcal{R}$ be a finite set of relations between $\gamma_1, \dots, \gamma_m$ such that

$$\pi_1(Y - \mathcal{K}, y) \simeq \langle \gamma_1, \dots, \gamma_m \mid \mathcal{R} \rangle$$

(fundamental groups of hypersurface complements are always finitely presentable; a way to prove this is by intersecting with a generic 2-plane and applying Van Kampen method). Choose actual paths representing the generators and choose homotopies realising the relations. All these homotopies occur within a compact subspace $C \subseteq Y$. The image $LL(C)$ is compact. Choose x a complex number with sufficiently large module, such that the horizontal section $\{x\} \times C$ does not contain any point of $\mathcal{H}$. Let $\tilde{\gamma}_1, \dots, \tilde{\gamma}_m$ be the elements of $\pi_1(E, (x, y))$ obtained by lifting in $\{x\} \times C$ the paths representing $\gamma_1, \dots, \gamma_m$. These elements satisfy the same defining relations (where each γ_j is replaced by $\tilde{\gamma}_j$ – we denote this set of relations by $\tilde{\mathcal{R}}$). The map $\gamma_j \mapsto \tilde{\gamma}_j$ is a section of p_* .

The fiber F identifies with $\mathbb{C} - LL(y)$, that is, since $y \notin \mathcal{K}$, the complement of n points in $\mathbb{C}$. Its fundamental group is a free group of rank n . A standard generating system $f_1, \dots, f_n$ consisting of generators-of-the-monodromy is obtained when choosing a *planar spider* (in the sequel, we will use a simpler way of encoding standard generators, by means of *tunnels*).

The group $\pi_1(E, (x, y))$ is generated by $f_1, \dots, f_n, \tilde{\gamma}_1, \dots, \tilde{\gamma}_m$. Each γ_j defines a monodromy automorphism $\phi_j \in \text{Aut}(\pi_1(F, (x, y)))$ such that, for all $i \in \{1, \dots, n\}$ and all $j \in \{1, \dots, m\}$, one has

$$\tilde{\gamma}_j f_i \tilde{\gamma}_j^{-1} = \phi_j(f_i).$$

The semi-direct product structure implies that we have a presentation

$$\pi_1(E, (x, y)) \simeq \langle f_1, \dots, f_n, \tilde{\gamma}_1, \dots, \tilde{\gamma}_m \mid \tilde{\gamma}_j f_i \tilde{\gamma}_j^{-1} = \phi_j(f_i), \tilde{\mathcal{R}} \rangle.$$

Since $\ker \beta$ is the normal subgroup generated the $\tilde{\gamma}_j$'s, we obtain

$$\pi_1(W \setminus \setminus V^{\text{reg}}, (x, y)) \simeq \langle f_1, \dots, f_n, \tilde{\gamma}_1, \dots, \tilde{\gamma}_m \mid \tilde{\gamma}_j f_i \tilde{\gamma}_j^{-1} = \phi_j(f_i), \tilde{\mathcal{R}}, \tilde{\gamma}_1 = 1, \dots, \tilde{\gamma}_m = 1 \rangle.$$

Since the relations in $\tilde{\mathcal{R}}$ do not involve $f_1, \dots, f_n$, the presentation may be rewritten as:

Theorem 4.8. *Let $\phi_1, \dots, \phi_m$ be monodromy automorphisms of the fundamental group of $L_y \cap W \setminus \setminus V^{\text{reg}}$ associated with generators $\gamma_1, \dots, \gamma_m$ of the fundamental group of $Y - \mathcal{K}$. The braid group of W admits the presentation*

$$\pi_1(W \setminus \setminus V^{\text{reg}}, (x, y)) \simeq \langle f_1, \dots, f_n \mid f_i = \phi_j(f_i) \rangle,$$

where one takes a relation $f_i = \phi_j(f_i)$ for each pair i, j with $i \in \{1, \dots, n\}$ and $j \in \{1, \dots, m\}$.

The above theorem refines the main result from [2], which unfortunately was written without knowledge of classical article by Looijenga [25].


Later on, we will be able to write presentations of $B(W)$ in much more explicit ways.

5. TUNNELS, LABELS AND HURWITZ RULE

Let W be an irreducible well-generated complex reflection group. We keep the notations from the previous section. Let $y \in Y$. Let U_y be the complement in L_y of the vertical imaginary half-lines below the points of $LL(y)$, or in more formal terms:

$$U_y := \{(z, y) \in L_y \mid \forall x \in LL(y), \text{re}(z) = \text{re}(x) \Rightarrow \text{im}(z) > \text{im}(x)\}.$$

In this example, the support of $LL(y)$ consists of 4 points, and U_y is the complement of three half-lines:



Let

$$\mathcal{U} := \bigcup_{y \in Y} U_y.$$

Lemma 5.1. *The subset $\mathcal{U}$ is dense in $W \setminus \setminus V^{\text{reg}}$, open and contractible.*

Proof. The first two statements are clear.

Define a continuous function $\beta : Y \rightarrow \mathbb{R}$ by

$$\beta(y) := \max\{\text{im}(x) \mid x \in LL(y)\} - 1.$$

Points of $W \setminus \setminus V$ are represented by pairs $(x, y) \in \mathbb{C} \times Y$, or equivalently by triples $(a, b, y) \in \mathbb{R} \times \mathbb{R} \times Y$, where $a = \text{re}(x)$ and $b = \text{im}(x)$. For $t \in [0, 1]$, define $\phi_t : W \setminus \setminus V \rightarrow W \setminus \setminus V$ by

$$\phi_t(a, b, y) := \begin{cases} (a, b, y) & \text{if } b \geq \beta(y), \\ (a, b + t(\beta(y) - b), y) & \text{if } b \leq \beta(y). \end{cases}$$

Each ϕ_t preserves $\mathcal{U}$ and the homotopy ϕ restricts to a deformation retraction of $\mathcal{U}$ to

$$\bigcup_{y \in Y} \{(x, y) \in L_y \mid \text{im}(x) \geq \beta(y)\}.$$


The latter is a locally trivial bundle over the contractible space Y , with contractible fibers (the fibers are half-planes). Thus it is contractible. $\square$

As explained in the topological preliminaries, we may (and will) use $\mathcal{U}$ as “basepoint” for $W \setminus V^{\text{reg}}$.

Definition 5.2. The *braid group* of W is $B(W) := \pi_1(W \setminus V^{\text{reg}}, \mathcal{U})$.

As for other notions actually depending on W , we usually write B instead of $B(W)$, since most of the time we implicitly refer to a given W .

Definition 5.3. A *semitunnel* is a triple $T = (y, z, L) \in Y \times \mathbb{C} \times \mathbb{R}_{\geq 0}$ such that $(z, y) \in \mathcal{U}$ and the affine segment $[(z, y), (z + L, y)]$ lies in $W \setminus V^{\text{reg}}$. The path γ_T associated with T is the path $t \mapsto (z + tL, y)$. The semitunnel T is a *tunnel* if in addition $(z + L, y) \in \mathcal{U}$.



The distinction between tunnels and semitunnels should be understood in light of our topological conventions: if T is a tunnel, γ_T represents an element

$$b_T \in \pi_1(W \setminus V^{\text{reg}}, \mathcal{U}),$$

while semitunnels will be used to represent points of the universal cover $(\widetilde{W \setminus V^{\text{reg}}})_{\mathcal{U}}$ (see Section 9).


Definition 5.4. An element $b \in B$ is *simple* if $b = b_T$ for some tunnel T .

Each tunnel lives in a given L_y , where it may be represented by an horizontal (= constant imaginary part) segment avoiding $LL(y)$ and with endpoints in U_y . The triple (y, z, L) may be uniquely recovered from $[(z, y), (z + L, y)]$. A frequent abuse of terminology will consist of using the term *tunnel* (or *semitunnel*) to designate either the triple (y, z, L) , or the segment $[(z, y), (z + L, y)]$, or the pair $(y, [z, z + L])$, depending on the context (in particular, when intersecting tunnels with geometric objects, the tunnels should be understood as affine segments).


Let $y \in Y$. Let $(x_1, \dots, x_k)$ be the ordered support of $LL(y)$. The space $p_x((L_y \cap W \setminus V^{\text{reg}}) - U_y)$ is an union of m disjoint open affine intervals $I_1, \dots, I_k$, where

$$I_i := \begin{cases} (x_i - \sqrt{-1}\infty, x_i) & \text{if } i = 1 \text{ or } (i > 1 \text{ and } \text{re}(x_{i-1}) < \text{re}(x_i)), \\ (x_{i-1}, x_i) & \text{otherwise.} \end{cases}$$

(by $(x_i - \sqrt{-1}\infty, x_i)$, we mean the open vertical half-line below x_i). In the first case (when I_i is not bounded), we say that x_i is *deep*.



Choose a *system of elementary tunnels* for y . By this, we mean the choice, for each $i = 1, \dots, k$, of a small tunnel T_i in L_y crossing I_i and not crossing the other intervals; let $s_i := b_{T_i}$ be the associated element of B .



These elements depend only on y and not of the explicit choice of elementary tunnels.

Definition 5.5. The sequence $lbl(y) := (s_1, \dots, s_k)$ is the *label* of y . Let $i_1, i_2, \dots, i_l$ be the indices of the successive deep points of $LL(y)$. The *deep label* of y is the subsequence $(s_{i_1}, \dots, s_{i_l})$.

In the above example, the deep label is (s_1, s_3, s_4) . The length of the label is n if and only if $y \in Y - \mathcal{K}$. The deep label coincides with the label if and only if $y \in Y^{\text{gen}}$.

It is clear that if $y \in Y^{\text{gen}}$, the elements $s_1, \dots, s_n$ such that $(s_1, \dots, s_n) = lbl(y)$ are braid reflections.

Later on, it will appear that the pair $(LL(y), lbl(y))$ uniquely determines y (Theorem 6.6). It is also possible to characterise, for a given configuration (not necessarily regular), the possible labels, thus to explicitly describe of the fibers of LL .

Consider the case $y = 0$ (given by the equations $X_1 = 0, \dots, X_{n-1} = 0$). The multiset $LL(y)$ has support $\{0\}$ with multiplicity n .

Definition 5.6. We denote by δ the simple element such that $lbl(0) = (\delta)$.

This element plays the role Deligne's element Δ . Let $v \in V^{\text{reg}}$ such that the W -orbit $\bar{v}$ lies in L_0 . Broué-Malle-Rouquier consider the element (denoted by π , [14, Notation 2.3]) in the pure braid group $P(W) := \pi_1(V^{\text{reg}}) = \ker(B(W) \rightarrow W)$ represented by the loop

$$t \mapsto v \exp(2\sqrt{-1}\pi t).$$

They observed that this element lies in the center of B ([14, Theorem 2.24]). Since X_n has weight h , δ^h coincides with this element π . More precisely, δ is represented by the loop in $W \setminus V^{\text{reg}}$ image of the path in V^{reg}

$$t \mapsto v \exp(2\sqrt{-1}\pi t/h).$$

In particular:

Lemma 5.7. *The element δ^h is central in B . The image of δ in W is ζ_h -regular, in the sense of [34].*

Each tunnel lives in a single fiber L_y . Simple elements represented by tunnels in different fibers may be compared since being a tunnel is an open condition, which is preserved when y is perturbed.

Definition 5.8. Let $T = (y, z, L)$ be a tunnel. A T -neighbourhood of y is a path-connected neighbourhood Ω of y in Y such that, for all $y' \in \Omega$, $T' := (y', z, L)$ is a tunnel.

Such neighbourhoods clearly exist for all $y \in Y$.

Lemma 5.9 (Hurwitz rule). *Let $T = (y, z, L)$ be a tunnel, representing a simple element s . Let Ω be a T -neighbourhood of y . For all $y' \in \Omega$, $T' := (y', z, L)$ represents s .*


Proof. This simply expresses that the tunnels (y', z, L) and (y, z, L) represent homotopic paths, which is clear by definition of Ω . $\square$

Remark 5.10. Let $T_1, \dots, T_k$ is a system of elementary tunnels for y . Let Ω_i be a T_i -neighbourhood for y . A *standard neighbourhood* of y could be defined as a path-connected neighbourhood Ω of y inside $\cap_{i=1}^k \Omega_i$. These standard neighbourhoods form a basis for the topology of Y . A consequence of Hurwitz rule is that the label of y may uniquely be recovered once we know the label of a single $y' \in \Omega$. The converse is not true, unless $y \in Y - \mathcal{K}$. Let $y' \in \Omega$, let S and S' be the strata of $LL(y)$ and $LL(y')$ (with respect to the discriminant stratification of E_n). We have $S' \leq S$. The label of y' is uniquely determined by the label of y if and only if $S = S'$. Otherwise, the ambiguity is caused by the extra ramification at y .

The remainder of this section consists of various consequences of Hurwitz rule.

Corollary 5.11. *Let $y \in Y$. Let $(s_{i_1}, \dots, s_{i_l})$ be the deep label of y . We have $s_{i_1} \dots s_{i_l} = \delta$.*

Proof. Any tunnel T deep enough and long enough represents $s_{i_1} \dots s_{i_l}$.




The origin $0 \in Y$ lies in a T -neighbourhood of y . To conclude, apply Hurwitz rule. $\square$

Corollary 5.12. *Let $x \in E_n^{\text{reg}}$. Let $\beta \in \pi_1(E_n^{\text{reg}}, x)$, $y \in LL^{-1}(x)$ and $y' := y \cdot \beta$. Let $(b_1, \dots, b_n)$ be the label of y and $(b'_1, \dots, b'_n)$ be the label of y' . Then*

$$(b'_1, \dots, b'_n) = (b_1, \dots, b_n) \cdot \beta,$$

where β acts by right Hurwitz action.

Proof. It is enough to prove this for a standard generator σ_i . Let $(x_1, \dots, x_n)$ be the ordered support of x . By Hurwitz rule, we may adjust the imaginary parts of the x_i 's without affecting the label; in particular we may assume that $\text{im}(x_i) < \text{im}(x_{i+1})$. We may find tunnels $T_- = (y, z_-, L)$ and $T_+ = (y, z_+, L)$ as in the picture below:



The path in E_n^{reg} where x_{i+1} moves along the dotted arrow and all other points are fixed represents σ_i . Applying Hurwitz rule to T_+ , we obtain $b'_i = b_{i+1}$; applying Hurwitz rule to T_- , we obtain $b'_i b'_{i+1} = b_i b_{i+1}$. The result follows. $\square$

Corollary 5.13. *Let $y \in Y^{\text{gen}}$. The cardinal of the Hurwitz orbit $\text{lbl}(y) \cdot B_n$ is at most $n!h^n/|W|$, and there is an equivalence between:*


- (i) $|\text{lbl}(y) \cdot B_n| = n!h^n/|W|$.
- (ii) *The map $Y^{\text{gen}} \rightarrow E_n \times B^n, y \mapsto (LL(y), \text{lbl}(y))$ is injective.*

In the next section, we will prove that both (i) and (ii) hold. This is not a trivial statement.

Proof. By the previous corollary, one has $\text{lbl}(y \cdot \beta) = \text{lbl}(y) \cdot \beta$. Since $y \in Y - \mathcal{K}$, we have $|y \cdot \beta| = n!h^n/|W|$ (Theorem 4.4). The corollary follows. $\square$

Corollary 5.14. *Let s be a simple element. There exists $y \in Y^{\text{gen}}$ and $i \in \{1, \dots, n\}$ such that $s = s_1 \dots s_i$, where $(s_1, \dots, s_n) := \text{lbl}(y)$.*

Proof. Let T be a tunnel representing s . Any T -neighbourhood of y contains generic points. We may assume that $y \in Y^{\text{gen}}$. The picture below explains, on an example, how to move certain points (following the dotted paths) of the underlying configuration to reach a suitable y' :



This path in E_n^{reg} lifts, via LL , to a path in a T -neighbourhood of y whose final point y' satisfies the conditions of the lemma. $\square$

6. REDUCED DECOMPOSITIONS OF COXETER ELEMENTS

Most results on this section follow from the generalisation to well-generated complex 2-reflection groups of a property of finite Coxeter groups initially conjectured by Looijenga, [25, (3.5)], and solved in a letter from Deligne to Looijenga (citing discussions with Tits and Zagier), [21]. This property is more or less equivalent to [3, Fact 2.2.4], used in the our earlier construction of the dual braid monoid. It relies on a case-by-case argument, which we hope to improve in the future.

As in the previous sections, W is a well-generated complex 2-reflection group, but not necessarily irreducible. We study the images of simple elements under the natural projection $\pi : B \rightarrow W$. Recall that the projection π is part of the fibration exact sequence

$$1 \longrightarrow \pi_1(V^{\text{reg}}) \longrightarrow \pi_1(W \setminus V^{\text{reg}}) \xrightarrow{\pi} W \longrightarrow 1.$$

For this exact sequence to be well-defined, one has to make consistent choices of basepoints in V^{reg} and in $W \setminus V^{\text{reg}}$. We have already described our “basepoint” $\mathcal{U}$ in $W \setminus V^{\text{reg}}$. Choose $u \in \mathcal{U}$ and choose a preimage $\tilde{u}$ of u in V^{reg} . If $u' \in \mathcal{U}$ is another choice and if γ is a path in $\mathcal{U}$ from u to u' , γ lifts to a unique path $\tilde{\gamma}$ starting at $\tilde{u}$; since $\mathcal{U}$ is contractible, the fixed-endpoint homotopy class of $\tilde{\gamma}$ (and, in particular, its final point) does not depend on γ . In other words, once we have chosen a preimage of one point of $\mathcal{U}$, we have a natural section $\tilde{\mathcal{U}}$ of $\mathcal{U}$ in V^{reg} , as well as a transitive system of isomorphisms between $(\pi_1(V^{\text{reg}}, \tilde{u}))_{\tilde{u} \in \tilde{\mathcal{U}}}$. From now on, we assume we have made such a choice, and we define the pure braid group as $\pi_1(V^{\text{reg}}, \tilde{\mathcal{U}})$. (The $|W|$ possible choices yield conjugate morphisms π).

Recall that an element $w \in W$ is ζ -regular, in the sense of [34], if it admits in ζ -eigenvector in V^{reg} . We then say that ζ is a *regular eigenvalue*, and its order d is a *regular number*. Springer proved that, if ζ is a regular eigenvalue, all ζ -regular elements of W are conjugate in W .

Definition 6.1. Let W be an irreducible well-generated complex 2-reflection group. An element $c \in W$ is a (*generalised*) *Coxeter element* if it is ζ_h -regular. More generally, if W is a well-generated complex 2-reflection group decomposed as a sum $W = \bigoplus_i W_i$ of irreducible groups, a Coxeter element in W is a product $c = \prod_i c_i$ of Coxeter elements in each W_i .

A consequence of our description of δ (see previous section) is that $c := \pi(\delta)$ is a Coxeter element. The other Coxeter elements, which are conjugate to c , appear when considering other basepoints over $\mathcal{U}$.

We denote by R the set of all reflections in W . As in [3], for all $w \in W$, we denote by $\text{Red}_T(w)$ the set of *reduced R -decompositions* of w , that is minimal length sequences of elements of R with product w . Since R is closed under conjugacy, $\text{Red}_T(c)$ is stable under Hurwitz action. We also consider the length function $l_R : W \rightarrow \mathbb{Z}_{\geq 0}$, whose value at w is the common length of the elements of $\text{Red}_R(w)$. We also consider the two partial orderings of W defined as follows: for all $w, w' \in W$, we set

$$w \preceq_R w' :\Leftrightarrow l_R(w) + l_R(w^{-1}w') = l_R(w')$$

and

$$w' \succcurlyeq_R w :\Leftrightarrow l_R(w') + l_R(w'w^{-1}) = l_R(w).$$

Since R is invariant by conjugacy, we have $w \preceq_R w' \Leftrightarrow w' \succcurlyeq_R w$.

It is worth noting that the basic Lemma 1.2.1. of [3] does not hold when W is not a complexified real group: for obvious reasons, one has $l_R(w) \geq n$ for all $w \in W$, but sometimes $l_R(w) > n$. However, we have:

Lemma 6.2. *Let c be a Coxeter element in a well-generated complex 2-reflection group W . Let $w \in W$ with $w \preceq_R c$. Then*

- $\text{codim ker}(w - 1) = l_R(w)$,
- w is a Coxeter element in the group $W_w := \{w' \in W \mid \ker(w - 1) \subseteq \ker(w' - 1)\}$,
- $\text{Red}_R(w) = \text{Red}_{R \cap W_w}(w)$.

That W_w is a reflection group is classical (the fixator of any subspace of V is a reflection group, called *parabolic subgroup*). One checks (e.g., on the classification) that parabolic subgroups of well-generated complex reflection groups are well-generated.

Proof. The lemma reduces to the case when W is irreducible, which we assume from now on.

(i) Assume that $w = c$. We have $\ker(c - 1) = 0$ since the eigenvalues of c are $\zeta_h^{1-d_1}, \dots, \zeta_h^{1-d_n}$ and $1 < d_i \leq h$ for all i . That $l_R(c) = n$ is easy and classical (it follows, for example, from the main theorem in [2] – in this reference, the statements about the eigenvalues contain a typo: ζ should be replaced by $\bar{\zeta}$).

(ii) Let $w \prec c$. Let us check that w is a Coxeter element in W_w . When W is complexified real, this follows from [3, Lemma 1.4.3]. For $G(e, e, r)$, it follows from [5, Proposition 4.3].

The remaining exceptional cases are easy (an induction reduces the problem to the case $l_R(w) = n - 1$) to check (though unfortunately we don't have a general argument).

(iii) We have $w = c_1 \dots c_k$, where the c_i are Coxeter elements of irreducible parabolic subgroups. By (i), we have, for each c_i , $l_{R_{c_i}}(c_i) = \text{codim ker}(c_i - 1)$. We deduce that $l_R(w) \leq \sum \text{codim ker}(c_i - 1) = \text{codim ker}(w - 1)$. The opposed inequality being obvious, we have $l_R(w) = \text{codim ker}(w - 1)$.

(iv) If $w = r_1 \dots r_k$ with $r_i \in R$ and $k = \text{codim ker}(w - 1)$, one necessarily has $\text{ker}(w - 1) = \bigcap_{i=1}^k \text{ker}(r_i - 1)$. In other words, $r_i \in W_w$. This proves the last statement. $\square$

Proposition 6.3. *Let W be a well-generated complex 2-reflection group. Let c be a generalised Coxeter element in W . Let $w \in W$ with $w \preceq c$. The set $\text{Red}_R(w)$ is a single Hurwitz orbit. If W is irreducible with Coxeter number h , we have $|\text{Red}_R(c)| = n!h^n/|W|$.*

Proof. Using Lemma 6.2, the proposition reduces to the case when W is irreducible and $w = c$: in the reducible case, reduced decompositions of Coxeter elements are “shuffles” of reduced decompositions of the Coxeter elements of the irreducible summands.

The complexified real case is studied in [21] (transitivity is easy and does not require case-by-case, see for example [3], Proposition 1.6.1). The $G(e, e, r)$ case is a combination of Proposition 6.1 (transitivity) and Theorem 8.1 (cardinality) from [5]. The remaining exceptional types are done by computer. $\square$

We now assume that W is an irreducible well-generated complex 2-reflection group and return to the situation and notations of the previous sections.

Theorem 6.4. *Let $y \in Y^{\text{gen}}$, with label $(s_1, \dots, s_n)$. For $i = 1, \dots, n$, let $r_i := \pi(s_i)$. The map π^n induces an isomorphism of B_n -sets between the Hurwitz orbits of $(s_1, \dots, s_n)$ and of $(r_1, \dots, r_n)$.*

Proof. We have $\delta = s_1 \dots s_n$. Let $c := r_1 \dots r_n$. As mentioned above, c is a Coxeter element ([2, Lemma 3.1]). By Proposition 6.3, the cardinal of the Hurwitz $(r_1, \dots, r_n) \cdot B_n$ is $n!h^n/|W|$. By Corollary 5.13, the cardinal of the Hurwitz orbit $(s_1, \dots, s_n) \cdot B_n$ is at most $n!h^n/|W|$. One concludes that $|(s_1, \dots, s_n) \cdot B_n| = n!h^n/|W|$ and that $\pi^n : (s_1, \dots, s_n) \cdot B_n \rightarrow (r_1, \dots, r_n) \cdot B_n$ is an isomorphism. $\square$

Corollary 6.5. *The restriction of π to the set S of simple elements is injective.*

Proof. Let s and s' be simples such that $\pi(s) = \pi(s')$. By Corollary 5.14, we may find $y, y' \in Y^{\text{gen}}$, with $\text{lbl}(y) = (s_1, \dots, s_n)$ and $\text{lbl}(y') = (s'_1, \dots, s'_n)$, and $i \in \{0, \dots, n\}$ such that $s = s_1 \dots s_i$ and $s' = s'_1 \dots s'_i$. By Proposition 6.3 applied to $w = \pi(s)$, we conclude that we may find $\beta \in B_i$ such that $(\pi(s_1), \dots, \pi(s_i)) \cdot \beta = (\pi(s'_1), \dots, \pi(s'_i))$. Similarly, we find $\beta' \in B_{n-i}$ such that $(\pi(s_{i+1}), \dots, \pi(s_n)) \cdot \beta' = (\pi(s'_{i+1}), \dots, \pi(s'_n))$. View $B_i \times B_{n-i}$ as a subgroup of B_n (the first factor braids the first i strings, the second factors braids the $n - i$ last strings), and set $\beta'' := (\beta, \beta') \in B_n$. We have $\pi^n(\text{lbl}(y)) \cdot \beta'' = \pi^n(\text{lbl}(y'))$. Applying the theorem, this implies that $\text{lbl}(y) \cdot \beta'' = \text{lbl}(y')$. Clearly, β'' does not modify the product of the first i terms of the labels. Thus $s = s'$. $\square$

In the following theorem, B^* denotes the set of finite sequences of elements of B .

Theorem 6.6. *The map $Y \rightarrow E_n \times B^*, y \mapsto (LL(y), \text{lbl}(y))$ is injective.*

The image of the map may easily be described in terms of simple axioms relating $lbl(y)$ to $LL(y)$. This description is not needed here.

Proof. The injectivity on Y^{gen} follows from Corollary 5.13 and the proof of the previous theorem.

Let $y, y' \in Y$ with $LL(y) = LL(y')$ and $lbl(y) = lbl(y')$. We have to prove that $y = y'$. Let $(x_1, \dots, x_k)$ and $(n_1, \dots, n_k)$ be the ordered support and multiplicities of $LL(y)$. Let $(t_1, \dots, t_k) := lbl(y)$.

For simplicity, we begin with the case when $\{x_1, \dots, x_k\} \in E_k^{\text{gen}}$, *i.e.*, we assume that $\text{re}(x_1) < \text{re}(x_2) < \dots < \text{re}(x_k)$. Let $(y_m)_{m \geq 0}$ be sequences of points in Y^{gen} converging to y . Let $(s_1^{(m)}, \dots, s_n^{(m)}) := lbl(y_m)$. For m large enough, Hurwitz rule implies that $s_1^{(m)} \dots s_{n_1}^{(m)} = t_1$, $s_{n_1+1}^{(m)} \dots s_{n_1+n_2}^{(m)} = t_2$, etc... Similarly, choose a sequence (y'_m) of points in Y^{gen} converging to y' ; one has the corresponding properties for $(s_1^{(m)'}, \dots, s_n^{(m)'}) := lbl(y'_m)$. For m large, $(s_1^{(m)}, \dots, s_{n_1}^{(m)})$ and $(s_1^{(m)'}, \dots, s_{n_1}^{(m)'})$ are two reduced R -decompositions of the simple t_1 . By Proposition 6.3, they are in the same B_{n_1} Hurwitz orbit. Similarly, $(s_{n_1+1}^{(m)}, \dots, s_{n_1+n_2}^{(m)})$ and $(s_{n_1+1}^{(m)'}, \dots, s_{n_1+n_2}^{(m)'})$ are in the same B_{n_2} Hurwitz orbit, etc... We have proved that, for m large enough, $lbl(y_m)$ and $lbl(y'_m)$ are Hurwitz transformed by a braid β_m which lies in the natural subgroup $B_{n_1} \times B_{n_2} \times \dots \times B_{n_k}$ of B_n . One concludes noting that this subgroup belongs to the ramification group for the B_n -action on y .

The case when $\{x_1, \dots, x_k\} \notin E_k^{\text{gen}}$ may be reduced to the previous case by some suitable perturbation. $\square$

We will also need the following lemma:

Lemma 6.7. *Let $y \in Y$. Let $T = (y, z, L)$ and $T' = (y, z', L')$ be two tunnels in L_y such that $b_T = b_{T'}$. Then T and T' cross the same intervals among $I_1, \dots, I_n$ (in other words, T and T' are homotopic as tunnels drawn in L_y).*

Proof. Up to perturbing y , we may assume that $y \in Y^{\text{gen}}$. Let $(s_1, \dots, s_n) := lbl(y)$, and $(r_1, \dots, r_n) := \pi^n(lbl(y))$. Let $i_1, \dots, i_l$ (resp. $j_1, \dots, j_m$) be the successive indices of the intervals among $I_1, \dots, I_n$ crossed by T (resp. T'). We have $b_T = s_{i_1} \dots s_{i_l}$ and $b_{T'} = s_{j_1} \dots s_{j_m}$. Assuming that $b_T = b_{T'}$, we obtain $s_{i_1} \dots s_{i_l} = s_{j_1} \dots s_{j_m}$ and $r_{i_1} \dots r_{i_l} = r_{j_1} \dots r_{j_m}$. Let $w := r_{i_1} \dots r_{i_l}$. By Lemma 6.2, $l = m$ and we have

$$(*) \quad \ker(w - 1) = \bigcap_{k=1}^l \ker(r_{i_k} - 1) = \bigcap_{k=1}^l \ker(r_{j_k} - 1).$$

Assume that $(i_1, \dots, i_l) \neq (j_1, \dots, j_l)$. We may find $j \in \{j_1, \dots, j_l\}$ such that, for example, $i_1 < \dots < i_k < j < i_{k+1} < \dots < i_l$. Noting that the element $r_{i_1} \dots r_{i_k} r_j r_{i_{k+1}} \dots r_{i_l}$ is a parabolic Coxeter element, we deduce that $\ker(r_j - 1) \not\subseteq \ker(w - 1)$. This contradicts $(*)$. $\square$

7. THE DUAL BRAID MONOID

Here again, W is an irreducible well-generated complex 2-reflection groups. We keep the notations from the previous sections. Recall that an element $b \in B$ is *simple* if $b = b_T$ for some tunnel T , and that the set of simple elements is denoted by S .

Definition 7.1. The *dual braid monoid* is the submonoid M of B generated by S .

Consider the binary relation $\preceq$ defined on S as follows. Let s and s' be simple elements. We write $s \preceq s'$ if and only if there exists $(z, y) \in W \setminus V^{\text{reg}}$, $L, L' \in \mathbb{R}_{\geq 0}$ with $L \leq L'$ such that (y, z, L) is a tunnel representing s and (y, z, L') is a tunnel representing s' .

This section is devoted to the proof of:

Theorem 7.2. *The monoid M is a Garside monoid, with set of simples S and Garside element δ . The relation $\preceq$ defined above on S is the restriction to S of the left divisibility order in M .*

The monoid M generates B , which inherits a structure of Garside group.

Several results mentioned in the introduction follow from this theorem:

Theorem 0.3 does not assume irreducibility, but follows immediately from the irreducible case since direct products of Garside groups are Garside groups.

Theorem 0.4 also follows, with arguments similar to [20, p. 301]: the center of B is generated by δ^a , where a is the smallest positive integer such that c^a is central in W (using standard techniques, one may also prove that all h -th roots of π in B are conjugate to δ , and compute the centralisers of the powers of δ).

Another important consequence of Theorem 7.2 is that one may associate a practical contractible simplicial complex acted on by B . To explain this, we need some additional terminology, imitating [8] and [16]:

Definition 7.3. Let G be a group, let Σ be a subset of G . We denote by $\text{Flag}(G, \Sigma)$ the simplicial complex as follows:

- (0) the set of vertices is G ,
- (1) the 1-simplices are the pairs $\{g, h\}$ of distinct elements of G such that $g^{-1}h \in \Sigma$ or $h^{-1}g \in \Sigma$,
- (k) for all $k > 1$, a $(k+1)$ -tuple $\{g_0, \dots, g_k\}$ of distinct elements of G is a k -simplex if and only if $\{g_i, g_j\}$ is a 1-simplex for all $i \neq j$.

In other words, $\text{Flag}(G, \Sigma)$ is the only flag complex whose 1-skeleton is the Cayley graph of (G, Σ) .

The following result is due, in this formulation, to Charney-Meier-Whittlesey. It is inspired by the work of Bestvina, [8], on spherical type Artin monoid (which itself is based on the work of Deligne). It was also implicitly used in several articles by T. Brady (e.g. [10]).

Theorem 7.4 ([16]). *Let G be a Garside group with set of simples Σ . Then $\text{Flag}(G, \Sigma)$ is contractible.*

Corollary 7.5. *The simplicial complex $\text{Flag}(B, S)$ is contractible.*

The strategy of proof of Theorem 7.2 is very similar to the one in [5].

Lemma 7.6. *Let $s \in S$. We have $l(s) = l_R(\pi(s))$.*

Proposition 7.7. *Let $s, s' \in S$. The following statements are equivalent:*

- (i) $s \preceq s'$,
- (ii) $\exists s'' \in S, ss'' = s'$,

(iii) $\pi(s) \preceq_R \pi(s')$.

Proof. (i) $\Rightarrow$ (ii). Assume that $s \preceq s'$ and choose $(x, y) \in W \setminus V^{\text{reg}}$ and $L \leq L'$ such that (y, x, L) is a tunnel representing s and (y, x, L') is a tunnel representing s' . Then $(y, x + L, L' - L)$ is a tunnel representing $s'' \in S$ such that $ss'' = s'$.

(ii) $\Rightarrow$ (iii). The natural length function l is additive on B . Thus, under (ii), we have $l(s) + l(s'') = l(s')$. On the other hand, for all $\sigma \in S$, $l(\sigma) = l_R(\pi(\sigma))$. Thus $l_R(\pi(s)) + l_R(\pi(s'')) = l_R(\pi(s'))$. Since $\pi(s'') = \pi(s)^{-1}\pi(s')$, this implies that $\pi(s) \preceq_R \pi(s')$.

(iii) $\Rightarrow$ (i). We may find $y, y' \in Y^{\text{gen}}$, with $lbl(y) = (s_1, \dots, s_n)$ and $lbl(y') = (s'_1, \dots, s'_n)$ such that $s = s_1 s_2 \dots s_{l(s)}$ and $s' = s'_1 s'_2 \dots s'_{l(s')}$. Set $w := \pi(s)$, $w' := \pi(s')$ and, for $i = 1, \dots, n$, set $r_i := \pi(s_i)$ and $r'_i := \pi(s'_i)$. Assuming (iii), we may find $r''_1, \dots, r''_{l(s')-l(s)} \in R$ such that $r_1 \dots r_{l(s)} r''_1 \dots r''_{l(s')-l(s)} = r'_1 \dots r'_{l(s')}$. The sequences

$$(r_1, \dots, r_{l(s)}, r''_1, \dots, r''_{l(s')-l(s)})$$

and

$$(r'_1, \dots, r'_{l(s')})$$

both lie in $\text{Red}_R(w')$. Since $w' \preceq c$, both sequences lie in the same Hurwitz orbit (Proposition 6.3). Thus

$$(r'_1, \dots, r'_n)$$

and

$$(r_1, \dots, r_{l(s)}, r''_1, \dots, r''_{l(s')-l(s)}, r'_{l(s')+1}, \dots, r'_n)$$

are transformed one onto the other by Hurwitz action of a braid $\beta \in B_n$ only braiding the first $l(s')$ strands. The Hurwitz transformed of

$$lbl(y') = (s'_1, \dots, s'_n)$$

by β is the label

$$(s''_1, \dots, s''_n)$$

of some $y'' \in Y^{\text{gen}}$. Since the braid only involves the first $l(s')$ strands, $s''_1 \dots s''_{l(s')} = s'_1 \dots s'_{l(s')} = s'$. One has $\pi(s''_i) = r_i$ for $i = 1, \dots, l(s)$, thus $\pi(s''_1 \dots s''_{l(s)}) = \pi(s_1 \dots s_{l(s)})$. By Corollary 6.5, this implies $s''_1 \dots s''_{l(s)} = s_1 \dots s_{l(s)} = s$. For x with small enough real and imaginary parts, one may find real numbers L, L' with $0 < L \leq L'$ such that (y'', x, L) is a tunnel with associated simple $s''_1 \dots s''_{l(s)} = s$ and (y'', x, L') is a tunnel with associated simple $s''_1 \dots s''_{l(s')} = s'$. $\square$

Proposition 7.8. *The map π restricts to an isomorphism $(S, \preceq) \xrightarrow{\sim} ([1, c], \preceq_R)$. In particular, $\preceq$ is an order relation on S .*

Proof. The previous proposition, applied to $s' = \delta$, proves that $\pi(S) \subseteq [1, c]$. It also proves that π induces a morphism of sets with binary relations $(S, \preceq) \rightarrow ([1, c], \preceq_R)$. The injectivity is Corollary 6.5.

Surjectivity: Choose $y \in Y^{\text{gen}}$. Let $(s_1, \dots, s_n) := lbl(y)$. Let $r_i := \pi(s_i)$. We have $(r_1, \dots, r_n) \in \text{Red}_R(c)$. Let $w \in [1, c]$. We may find $(r'_1, \dots, r'_n) \in \text{Red}_R(c)$ such that $r'_1 \dots r'_{l_T(w)} = w$. By Proposition 6.3, $(r'_1, \dots, r'_n)$ is Hurwitz transformed of $(r_1, \dots, r_n)$, thus there exists $y' \in Y^{\text{gen}}$ such that $\pi_*(lbl(y')) = (r'_1, \dots, r'_n)$. The simple element which is the product of the first $l(w)$ terms of $lbl(y')$ is in $\pi^{-1}(w)$. $\square$

We have the key lemma:

Lemma 7.9. *The poset $([1, c], \preceq_R)$ is a lattice.*

Proof. The real case is done in [3]. The $G(e, e, r)$ is done in [5]. The remaining cases may be done by computer. $\square$

Definition 7.10. The *dual braid relations* with respect to W and c are all the formal relations of the form

$$rr' = r'r'',$$

where $r, r', r'' \in R$ are such that $r \neq r'$, $rr' \in [1, c]$, and the relation $rr' = r'r''$ holds in W .

Clearly, dual braid relations only involve reflections in $R \cap [1, c]$. When W is complexified real, $R \subseteq [1, c]$. This does not hold in general (see the tables at the end of the article).

Lemma 7.11. *Let $R_c := R \cap [1, c]$. The monoid $\mathbf{M}(P_c)$ admits the monoid presentation*

$$\mathbf{M}(P_c) \simeq \langle R_c \mid \text{dual braid relations} \rangle.$$

Remark 7.12. When viewed as a group presentation, the presentation of the lemma is a presentation for $\mathbf{G}(P_c)$. As soon as we prove Theorem 7.2, Lemma 7.11 will give an explicit presentation for B . A way to reprove Theorem 0.5 is by simplifying the (redundent) presentation given by the lemma. This does not involve any monodromy computation.

Proof of Lemma 7.11. By definition, $\mathbf{M}(P_c)$ admits the presentation with generators R_c and a relation $r_1 \dots r_k = r'_1 \dots r'_k$ for each pair $(r_1, \dots, r_k), (r'_1, \dots, r'_k)$ of reduced R -decompositions of the same element $w \in [1, c]$. Call these relations *Hurwitz relations*. By transitivity of the Hurwitz action on $\text{Red}_R(c)$ (Proposition 6.3), the Hurwitz relations are consequences of the dual braid relations. The dual braid relations clearly hold in $\mathbf{M}(P_c)$ (to see this, complete (r, r') to an element $(r, r', r_3, \dots, r_n) \in \text{Red}_R(c)$). This proves the lemma. $\square$

Proof of Theorem 7.2. Set $R_c := R \cap [1, c]$. We are in the situation of subsection 0.4 in [3], (W, R_c) is a *generated group* and c is *balanced* (one first observes that $\{w \mid w \preceq_{R_c} c\} = \{w \mid w \preceq_R c\}$ and $\{w \mid c \succ_{R_c} w\} = \{w \mid c \succ_R w\}$; one concludes noting that c is balanced with respect to (W, R) , which is immediate since R is invariant by conjugacy). By Lemma 7.9 and [3, Theorem 0.5.2], the premonoid $P_c := [1, c]$ (together with the natural partial product) is a Garside premonoid. We obtain a Garside monoid $\mathbf{M}(P_c)$.

By Proposition 7.8, the restriction of π is a bijection from S to P_c . Let ϕ be the inverse bijection. Let $w, w' \in P_c$. Assume that the product $w.w'$ is defined in P_c . Let w'' be the value of this product. One has $w \preceq_R w''$, thus $\phi(w) \preceq \phi(w'')$ (using again Proposition 7.8), and we may find b' in S such that $\phi(w)b' = \phi(w'')$ (Proposition 7.7). Claim: $b' = \phi(w')$. Indeed, b' and $\phi(w')$ are two simple elements whose image by π is $w^{-1}w''$; one concludes using Corollary 6.5. This proves that ϕ induces a premonoid morphism $P_c \rightarrow S$ (where S is equipped with the restriction of the monoid structure), thus induces a monoid morphism $\mathbf{M}(P_c) \rightarrow M$ and a group morphism $\Phi : \mathbf{G}(P_c) \rightarrow B$.

Let us prove that Φ is an isomorphism. Choose a basepoint $y \in Y^{\text{gen}}$. Let $\gamma_1, \dots, \gamma_n$ be generators of $\pi_1(Y - K, y)$. Let $(s_1, \dots, s_n)$ be the label of y . Let us reinterpret the

presentation from Theorem 4.8 in terms of Hurwitz action. Since $LL : Y - \mathcal{K} \rightarrow E_n^{\text{gen}}$ is a covering, $\pi_1(Y - \mathcal{K}, y)$ may be identified with a subgroup $H \subseteq B_n$. The generators $f_1, \dots, f_n$ in Theorem 4.8 may be chosen to be $s_1, \dots, s_n$, and the monodromy automorphism $\phi_1, \dots, \phi_m$ are obtained by Hurwitz action on $s_1, \dots, s_n$. Let $h \in H$. Let $(s'_1, \dots, s'_n) := h(s_1, \dots, s_n)$ (by this, we mean the Hurwitz action of h on the free group generated by $s_1, \dots, s_n$; the s'_i 's are words in the s_i 's). Call *Van Kampen relations* the relations of the type $s'_i = s_i$, for any $i \in \{1, \dots, n\}$, $h \in H$, and s'_i obtained as above. We have

$$B \simeq \langle s_1, \dots, s_n \mid \text{Van Kampen relations} \rangle.$$

The map π induces a bijection from $A := \pi^{-1}(R_c) \subseteq S$ to R_c . Let $r_1, \dots, r_n$ be the images of $s_1, \dots, s_n$. By transitivity of Hurwitz action on $\text{Red}_R(c)$, the group $\mathbf{G}(P_c)$ is generated by $r_1, \dots, r_n$, the remaining generators in the presentation of Lemma 7.11 appearing as conjugates of $r_1, \dots, r_n$ (by successive use of dual braid relations). Our generating sets are compatible and the morphism

$$\Phi : \mathbf{G}(P_c) \simeq \langle R_c \mid \text{dual braid relations} \rangle \rightarrow B \simeq \langle s_1, \dots, s_n \mid \text{Van Kampen relations} \rangle$$

is defined by $r_i \mapsto s_i$. Add to the presentation of B formal generators indexed by

$$A - \{s_1, \dots, s_n\}$$

as well as the dual braid relations $\pi(r)\pi(r') = \pi(r')\pi(r'')$. Since the relations already hold in $\mathbf{G}(P_c)$, they hold in B , and we obtain a new presentation

$$B \simeq \langle A \mid \text{Van Kampen relations on } \{s_1, \dots, s_n\}, \text{ dual braid relations on } A \rangle.$$

To conclude that Φ is an isomorphism, it is enough to observe that the dual braid relations encode the full Hurwitz action of B_n on $(s_1, \dots, s_n)$, while the Van Kampen relations encode the action of $H \subseteq B_n$: thus Van Kampen relations are consequences of dual braid relations, and $\mathbf{G}(P_c)$ and B are given by equivalent presentations.

Since Φ is an isomorphism and $\mathbf{M}(P_c)$ naturally embeds in $\mathbf{G}(P_c)$ (this is a crucial property of Garside monoids, [18]), $\mathbf{M}(P_c)$ is isomorphic to its image M in B . The rest of theorem is clear. $\square$

As mentioned earlier, one may view B as a “reflection group”, generated by the set $\mathcal{R}$ of all braid reflections. An element of M is in $\mathcal{R}$ if and only if it has length 1 for the natural length function, or equivalently if it is an *atom* (*i.e.*, an element which has no strict divisor in M except the unity). By Proposition 6.3 and Theorem 6.4, there is a bijection between $\text{Red}_R(c)$ and the image of $lbl : Y^{\text{gen}} \rightarrow \mathcal{R}^n$. This image is clearly contained in $\text{Red}_{\mathcal{R}}(\delta)$. The conjecture below claims an analog in B of the transitivity of the Hurwitz action on $\text{Red}_R(c)$ (δ is the natural substitute for a Coxeter element in B). It implies that any element in $\text{Red}_{\mathcal{R}}(\delta)$ is the label of some $y \in Y^{\text{gen}}$.

Conjecture 7.13. *The Hurwitz action of B_n on $\text{Red}_{\mathcal{R}}(\delta)$ is transitive.*

The analog of this holds in free groups.

8. CHAINS OF SIMPLES

Here again, W is an irreducible well-generated complex 2-reflection group, and the notations from the previous section are still in use. For $k = 0, \dots, n$, we denote by $\mathcal{C}_k$ the set of (strict) chains in $S - \{1\}$ of cardinal k , *i.e.*, the set of k -tuples $(c_1, \dots, c_k)$ in S^k such that

$$1 \prec c_1 \prec \dots \prec c_k,$$

or equivalently the set of k -tuples $(c_1, \dots, c_k)$ in M^k such that


$$1 \prec c_1 \prec \dots \prec c_k \preceq \delta.$$

It is convenient to write $\{1 \prec c_1 \prec \dots \prec c_k\}$ instead of $(c_1, \dots, c_k)$.

We set $\mathcal{C} := \bigsqcup_{k=0}^n \mathcal{C}_k$.

Let $C := \{1 \prec c_1 \prec \dots \prec c_k\} \in \mathcal{C}$. We say that $y \in Y$ *represents* C if there exists $x \in U_y$ and real numbers $L_1, \dots, L_k$ such that $0 < L_1 < \dots < L_k$ and, for $i = 1, \dots, k$, (y, x, L_i) is a tunnel representing C_i .

Example: if $LL(y)$ is as in the illustration below and $(s_1, \dots, s_5) = lbl(y)$, y represents $1 \prec s_1 \prec s_1 s_3 \prec s_1 s_3 s_4 \prec s_1 s_3 s_4 s_5$, $1 \prec s_2 \prec s_2 s_4 \prec s_2 s_4 s_5$, $1 \prec s_2 \prec s_2 s_5$ and their subchains, but does not represent $1 \prec s_2 \prec s_2 s_3$ nor $1 \prec s_3 \prec s_3 s_5$.



Definition 8.1. For all $C \in \mathcal{C}$, we set $Y_C := \{y \in Y \mid y \text{ represents } C\}$.

To illustrate this notion, we observe that implication (ii) $\Rightarrow$ (i) from Proposition 7.7 expresses that for all $C \in \mathcal{C}_2$, Y_C is non-empty. Based on the results from the previous sections, this easily generalises to:

Lemma 8.2. *For all $C \in \mathcal{C}$, the space Y_C is non-empty.*

The goal of this section is to prove:

Proposition 8.3. *For all $C \in \mathcal{C}$, the space Y_C is contractible.*

This technical result will be used in Section 9, when studying the nerve of an open covering of the universal cover of $W \backslash V^{\text{reg}}$: we will need to prove that certain non-empty intersections of open sets are contractible, and these intersections will appear as fiber bundles over some Y_C , with contractible fibers.

The proposition is not very deep nor difficult but somehow *unconvenient* to prove since the retraction will be described via LL , through ramification points. The following particular cases are easier to obtain:

- If C is the chain $1 \prec \delta$, then $Y_C = Y \simeq \mathbb{C}^{n-1}$.
- More significantly, let W be a complex reflection group of type A_2 . Up to renormalisation, the discriminant is $X_2^2 + X_1^3$. Identify Y with $\mathbb{C}$. For all $y \in \mathbb{C}$, $LL(y) = \{\pm(-y)^{3/2}\}$. In particular, $LL(1) = \{\pm\sqrt{-1}\}$. Let s be the simple element represented by the tunnel with $y = 1$, $x = -1$ and $L = 2$. Let C be the chain $1 \prec s$. Then Y_C is the open cone consisting of non-zero elements of $\mathbb{C}$ with argument in the open interval $(-2\pi/3, 2\pi/3)$.

- Assume that $C \in \mathcal{C}_n$. All points in Y_C are generic. Consider the map $Y^{\text{gen}} \rightarrow \mathcal{C}_n \times E_n$ sending y to the pair $(\{1 \prec s_1 \prec s_1 s_2 \prec \cdots \prec s_1 s_2 \cdots s_n = \delta\}, LL(y))$, where $(s_1, \dots, s_n) = \text{lbl}(y)$. This map is an homeomorphism. The $(Y_C)_{C \in \mathcal{C}_n}$ are the connected components of Y^{gen} . Each of these components is homeomorphic to E_n , which is contractible. These $(Y_C)_{C \in \mathcal{C}_n}$ are some analogs of chambers.

In the following proposition, if $A \subseteq LL(y)$ is a submultiset, the *deep label* of A is the sequence $(t_1, \dots, t_p)$ of labels (with respect to y) of points in the support of A which are deep with respect to A (since these points may not be deep in $LL(y)$, the deep label of A is not necessarily a subsequence of the deep label of y).

Proposition 8.4. *Let $C = \{1 \prec c_1 \prec c_2 \prec \cdots \prec c_m\}$ be a chain in $\mathcal{C}_m$. Let $y \in Y$, let $(x_1, \dots, x_k)$ be the ordered support of $LL(y)$ and $(s_1, \dots, s_k)$ be the label of y . The following assertions are equivalent:*

- (i) $y \in Y_C$.
- (ii) *There exists a partition of $LL(y)$ into $m + 1$ submultisets $A_0, \dots, A_m$ such that:*
 - (a) *For all $i, j \in \{1, \dots, m\}$ with $i < j$, for all $x \in A_i$ and all $x' \in A_j$, we have $\text{re}(x) < \text{re}(x')$.*
 - (b) *For all $x \in A_0$, one has*

$$\text{re}(x) < \min_{x' \in A_1 \cup \dots \cup A_m} \text{re}(x')$$

or

$$\text{im}(x) < \min_{x' \in A_1 \cup \dots \cup A_m} \text{im}(x')$$

or

$$\text{re}(x) > \max_{x' \in A_1 \cup \dots \cup A_m} \text{re}(x').$$


- (c) *For all $i \in \{1, \dots, m\}$, the product of the deep label of A_i is c_i .*

Moreover, in situation (ii), the partition $LL(y) = A_0 \sqcup \cdots \sqcup A_m$ is uniquely determined by y and C .

The picture below illustrates the proposition for particular y and C . Here $\text{lbl}(y) = (s_1, \dots, s_5)$ and the considered chain is

$$C = \{1 \prec s_2 \prec s_2 s_4\}.$$

We have chosen tunnels $T_1 = (y, x, L_1)$ and $T_2 = (y, x, L_2)$, with $L_1 < L_2$, such that $b_{T_1} = s_2$ and $b_{T_2} = s_2 s_4$. The dotted lines represent these tunnels, as well as the vertical half-lines above x , $x + L_1$ and $x + L_2$. They partition the complex line into three connected components; the partition $A_0 \sqcup A_1 \sqcup A_2$ is the associated partition of $LL(y)$. It is clear the possibility of drawing the tunnels is subject precisely to the conditions on $A_0 \sqcup A_1 \sqcup A_2$ expressed in the proposition.



Proof. (i) $\Rightarrow$ (ii): $(y, z, L_1), \dots, (y, z, L_m)$ be tunnels representing the successive non-trivial terms of $C = \{1 \prec c_1 \prec \dots \prec c_m\}$. Set $z_0 := z$ and, for $i = 1, \dots, m$, $z_i := z + L_i$. We have

$$\operatorname{re}(z_0) < \operatorname{re}(z_1) < \dots < \operatorname{re}(z_m)$$

and

$$\operatorname{im}(z_0) = \operatorname{im}(z_1) = \dots = \operatorname{im}(z_m).$$

For $i \in \{1, \dots, m\}$, let A_i be the submultiset of $LL(y)$ consisting of points x such that $\operatorname{re}(z_{i-1}) < \operatorname{re}(x) < \operatorname{re}(z_i)$ and $\operatorname{im}(x) > \operatorname{im}(z_0)$. Let A_0 be the complement in $LL(y)$ of $A_1 \cup \dots \cup A_m$. One easily checks (a), (b) and (c).

(ii) $\Rightarrow$ (i): Conversely, assume we are given a partition $A_0 \sqcup A_1 \sqcup \dots \sqcup A_m$ satisfying conditions (a) and (b). One may recover tunnels $(y, z, L_1), \dots, (y, z, L_m)$ such that the above construction yields the partition $A_0 \sqcup A_1 \sqcup \dots \sqcup A_m$. Condition (c) then implies that the tunnels represent the elements of C , thus that $y \in Y_C$.

Unicity of the partition: this follows from condition (c) and Lemma 6.7. $\square$

Lemma 8.5. *Let $C = \{1 \prec c_1 \prec c_2 \prec \dots \prec c_m\} \in \mathcal{C}$. Define Y_C^0 as the subspace of Y_C consisting of points y whose associated partition $A_0, \dots, A_m$ (from Proposition 8.4 (ii)) satisfies the following conditions:*

- for $i = 0, \dots, m$, the support of A_i is a singleton $\{a_i\}$ and,
- $\operatorname{re}(a_0) = \min_{i=1, \dots, m} \operatorname{re}(a_i) - 1$ and $\operatorname{im}(a_0) = \min_{i=1, \dots, m} \operatorname{im}(a_i) - 1$.

Then Y_C^0 is contractible.

Proof. Let $y \in Y_C^0$. The support $(x_0, \dots, x_m)$ of $LL(y)$ is generic, thus the label $(s_0, \dots, s_m)$ of y coincides with the deep label, and we have $s_0 \dots s_m = \delta$ (Corollary 5.11). By Proposition 8.4, Condition (ii)(c), we have, for $i = 1, \dots, m$, $s_i = c_i$. Thus $s_0 = \delta(c_1 \dots c_m)^{-1}$. We have proved that the label of any $y \in Y_C^0$ must be $(\delta(c_1 \dots c_m)^{-1}, c_1, \dots, c_m)$. A consequence of Theorem 6.6 is that the map $Y_C^0 \rightarrow E_m^{\text{gen}}$ sending y to $(x_1, \dots, x_m)$ is an homeomorphism. One concludes with Lemma 3.2. $\square$


Regular coverings satisfy unique homotopy lifting property. The map $LL : Y \rightarrow E_n$ is not regular on $\mathcal{K}$ but is *stratified regular*, where E_n is endowed with the natural stratification: at each $y \in Y$, the Jacobian of LL is surjective onto the tangent space of the stratum of $LL(y)$. The following lemma says that paths of non-decreasing ramification admits unique lifts. It is a general property of stratified regular maps, though we only need this particular example.

Lemma 8.6. *Let $\gamma : [0, 1] \rightarrow E_n$ be a path. For all $t \in [0, 1]$, denote by S_t the stratum of the natural stratification containing $\gamma(t)$. Assume that $\forall t, t' \in [0, 1], t \leq t' \Rightarrow S_t \leq S_{t'}$. Let $y_0 \in LL^{-1}(\gamma(0))$. There exists a unique path $\tilde{\gamma} : [0, 1] \rightarrow Y$ such that $\gamma = LL \circ \tilde{\gamma}$ and $\tilde{\gamma}(0) = y_0$.*

Proof. Left to the reader. $\square$

We may now proceed to the proof of Proposition 8.3. Let $C = \{1 \prec c_1 \prec \dots \prec c_m\} \in \mathcal{C}$. Let $y \in Y_C$. Let $(x_1, \dots, x_k)$ be the ordered support of $LL(y)$, and $(n_1, \dots, n_k)$ be the multiplicities. Let $A_0, \dots, A_m$ be the partition of $LL(y)$ described in Proposition 8.4 (ii).

The picture below gives an idea of the retraction of Y_C onto Y_C^0 that will be explicitly constructed. It illustrates the motion of a given point $y \in Y_C$; the black dots indicate the support of $LL(y)$ and the arrows how this support moves during the retraction.



For $i = 1, \dots, m$, consider the multiset mass centre

$$a_i := \frac{\sum_{x \in A_i} x}{|A_i|}$$

(in this expression, A_i is viewed as a multiset: each x_j in A_i is taken n_j times, and $|A_i|$ is the multiset cardinal, *i.e.*, the sum of the n_j such that $x_j \in A_i$).

For each $t \in [0, 1]$, let

$$\gamma_y(t) := A_0 \cup \bigcup_{i=1}^m \{(1-t)x + ta_i | x \in A_i\}$$

(here again, we consider the multiset union; in particular, the multicardinal of $\gamma_y(t)$ is constant, equal to n — *i.e.*, $\gamma_y(t) \in E_n$). This defines a path in E_n .

The path γ_y satisfies the hypothesis of Lemma 8.6 and uniquely lifts to a path $\tilde{\gamma}_y$ in Y such that $\tilde{\gamma}_y(0) = y$. An easy consequence of Proposition 8.4 is that $\tilde{\gamma}_y$ is actually drawn in Y_C .

Let $y' := \tilde{\gamma}_y(1)$. Let

$$R := \min_{i=1, \dots, m} \operatorname{re}(a_i) = \operatorname{re}(a_1)$$

and

$$I := \min_{i=1, \dots, m} \operatorname{im}(a_i).$$

For all $x \in A_0$, we have $\operatorname{re}(x) < R$ or $\operatorname{im}(x) < I$ or $\operatorname{re}(x) > \operatorname{re}(a_m)$; denote by x' the complex number with the same real part as x and imaginary part $I - 1$; let $a_0 := R - 1 + \sqrt{-1}(I - 1)$. Consider the path $\beta_x : [0, 1] \rightarrow \mathbb{C}$ defined by:

$$\beta_x(t) := \begin{cases} (1-2t)x + 2tx' & \text{if } t \leq 1/2 \\ (2-2t)x' + (2t-1)a_0 & \text{if } t \geq 1/2. \end{cases}$$

For all $t \in [0, 1]$, one has $\operatorname{re}(\beta_x(t)) < R$ or $\operatorname{im}(\beta_x(t)) < I$ or $\operatorname{re}(\beta_x(t)) > \operatorname{re}(a_m)$.

The path $\gamma'_y : [0, 1] \rightarrow E_n$ defined by

$$\gamma'_y(t) := \underbrace{\{a_1, \dots, a_1\}}_{|A_1| \text{ times}} \cup \dots \cup \underbrace{\{a_m, \dots, a_m\}}_{|A_m| \text{ times}} \cup \bigcup_{x \in A_0} \beta_x(t)$$

satisfies the hypothesis of Lemma 8.6 and lifts to a unique path $\tilde{\gamma}'_y$ in Y such that $\tilde{\gamma}'_y(0) = y'$. Once again, a direct application of Proposition 8.4 ensures that $\tilde{\gamma}'_y$ is actually in Y_C . The endpoint $y'' := \tilde{\gamma}'_y(1)$ lies in the subspace Y_C^0 of Corollary 8.5.

The map

$$\begin{aligned} \varphi : Y_C \times [0, 1] &\longrightarrow Y_C \\ (y, t) &\longmapsto (\tilde{\gamma}'_y \circ \tilde{\gamma}_y)(t) \end{aligned}$$

is a retraction of Y_C onto its contractible subspace Y_C^0 . Thus Y_C is contractible.

9. THE UNIVERSAL COVER OF $W \setminus\!\!\setminus V^{\text{reg}}$

The main result of this section is:

Theorem 9.1. *The universal cover of $W \setminus\!\!\setminus V^{\text{reg}}$ is homotopy equivalent to $\text{Flag}(B, S)$.*

Combined with Corollary 7.5, this proves our main result Theorem 0.2 in the irreducible case. The reducible case follows.

To prove this theorem, we construct an open covering $\widehat{\mathcal{U}} = (\widehat{\mathcal{U}}_b)_{b \in B}$, such that the nerve of $\mathcal{U}$ is $\text{Flag}(B, S)$ (Proposition 9.6) and intersections of elements of $\widehat{\mathcal{U}}$ are either empty or contractible (Proposition 9.7). Under these assumptions, a standard theorem from algebraic topology ([23, 4G.3]) gives the desired result.

As explained in the topological preliminaries, our “basepoint” $\mathcal{U}$ provides us with a model denoted $(\widehat{W \setminus\!\!\setminus V^{\text{reg}}})_{\mathcal{U}}$ or simply $\widehat{W \setminus\!\!\setminus V^{\text{reg}}}$ for the universal cover of $W \setminus\!\!\setminus V^{\text{reg}}$. Recall that, to any semitunnel T , one associates a path γ_T whose source is in $\mathcal{U}$, thus a point in $\widehat{W \setminus\!\!\setminus V^{\text{reg}}}$. Moreover, we have a left action of $B = \pi_1(W \setminus\!\!\setminus V^{\text{reg}}, \mathcal{U})$ on $\widehat{W \setminus\!\!\setminus V^{\text{reg}}}$. With these conventions, our open covering is very easy to define:

Definition 9.2. The set $\widehat{\mathcal{U}}_1$ is the subset of $\widehat{W \setminus\!\!\setminus V^{\text{reg}}}$ of elements represented by semitunnels. For all $b \in B$, we set $\widehat{\mathcal{U}}_b := b\widehat{\mathcal{U}}_1$.

It is clear that two semitunnels represent the same point in $\widehat{W \setminus\!\!\setminus V^{\text{reg}}}$ if and only if they are equivalent in the following sense:

Definition 9.3. Two semitunnels $T = (y, x, L)$ and $T' = (y', x', L')$ are *equivalent* if and only if $y = y'$, $x + L = x' + L'$ and the affine segment $[(x, y), (x', y)]$ is included in $\mathcal{U}$.

Let $T = (y, x, L)$ be a semitunnel. The point of $\widehat{\mathcal{U}}_1$ determined by T , or in other words the equivalence class of T , is uniquely determined by y , $x + L$ and

$$\lambda(T) := \inf \{ \lambda' \in [0, L] \mid [(x, y), (x + L - \lambda', y)] \subseteq \mathcal{U} \}.$$

The number λ is the infimum of the length of semitunnels in the equivalence class of T . This infimum is not a minimum (unless T is included in $\mathcal{U}$, since $(y, x + L - \lambda(T), \lambda(T))$ is not in general a semitunnel).

We consider the following subsets of $\widehat{\mathcal{U}}_1$:

- ($\mathcal{U}_1$) If $\lambda(T) = 0 = \min \{ \lambda' \in [0, L] \mid [x, x + L - \lambda'] \subseteq U_y \}$, then T is a tunnel, equivalent to $(y, x + L, 0)$. Elements of $\widehat{\mathcal{U}}_1$ represented by such tunnels of length 0 form an open subset denoted by $\mathcal{U}_1$. This subset is actually a sheet over $\mathcal{U}$ of the universal covering, corresponding to the trivial lift of the “basepoint”.
- ($\overline{\mathcal{U}}_1$) We denote by $\overline{\mathcal{U}}_1$ the subset consisting of points represented by tunnels with $\lambda(T) = 0$ (but without requiring that the “inf” is actually a “min”). We obviously have $\mathcal{U}_1 \subseteq \overline{\mathcal{U}}_1$, and $\overline{\mathcal{U}}_1$ is contained in the closure of $\mathcal{U}_1$.

Similarly, we set for all $b \in B$

$$\mathcal{U}_b := b\mathcal{U}_1, \quad \overline{\mathcal{U}}_b := b\overline{\mathcal{U}}_1.$$

Also, for any $y \in Y$, we denote by $\widehat{\mathcal{U}}_{b,y}$ (resp. $\mathcal{U}_{b,y}$, resp. $\overline{\mathcal{U}}_{b,y}$) the intersection of $\widehat{\mathcal{U}}_b$ (resp. $\mathcal{U}_b$, resp. $\overline{\mathcal{U}}_b$) with the fiber over y of the composed map $\widetilde{W \setminus V^{\text{reg}}} \rightarrow W \setminus V^{\text{reg}} \rightarrow Y$.

Lemma 9.4. *The family $(\overline{\mathcal{U}}_b)_{b \in B}$ is a partition of $\widetilde{W \setminus V^{\text{reg}}}$.*

Proof. It suffices to show that the projection $\overline{\mathcal{U}}_1 \rightarrow W \setminus V^{\text{reg}}$ is bijective. Any point in $z \in W \setminus V^{\text{reg}}$ is the target of a unique equivalence class of semitunnels T with $\lambda(T) = 0$; depending on whether $z \in \mathcal{U}$ or not, the associated point will be in $\mathcal{U}_1$ or in $\overline{\mathcal{U}}_1 - \mathcal{U}_1$. $\square$

In particular, $(\widehat{\mathcal{U}}_b)_{b \in B}$ is a covering of $\widetilde{W \setminus V^{\text{reg}}}$.

Lemma 9.5. *For all $b \in B$, $\widehat{\mathcal{U}}_b$ is open and contractible.*

Proof. It is enough to deal with $b = 1$. That $\widehat{\mathcal{U}}_1$ is open is easy. Let $\mathcal{T}$ be the space of tunnels and $\sim$ the equivalence relation. As a set, $\widehat{\mathcal{U}}_1 \simeq \mathcal{T} / \sim$. Consider the map

$$\begin{aligned} \phi : \mathcal{T} \times [0, 1] &\longrightarrow \mathcal{T} \\ (T = (y, x, L), t) &\longmapsto (y, x, L - t\lambda(T)) \end{aligned}$$

If $T \sim T'$, then $\forall t, \phi(T, t) \sim \phi(T', t)$. Thus ϕ induces a map $\overline{\phi} : \widehat{\mathcal{U}}_1 \times [0, 1] \rightarrow \widehat{\mathcal{U}}_1$. This map is continuous (this follows from the fact that λ induces a continuous function on $\widehat{\mathcal{U}}_1$) and $\overline{\phi}(T, t) = T$ if $T \in \overline{\mathcal{U}}_1$ or if $t = 0$. We have proved that $\widehat{\mathcal{U}}_1$ retracts to $\overline{\mathcal{U}}_1$. We are left with having to prove that $\overline{\mathcal{U}}_1$ is contractible. Inside $\overline{\mathcal{U}}_1$ lies $\mathcal{U}_1$ which is contractible, since it is a standard lift of the contractible space $\mathcal{U}$.

We conclude by observing that $\mathcal{U}_1$ and $\overline{\mathcal{U}}_1$ are homotopy equivalent. There is probably a standard theorem from semialgebraic geometry applicable here, but I was unable to find a proper reference. Below is a “bare-hand” argument: it explains how $\overline{\mathcal{U}}_1$ may be “locally retracted” inside $\mathcal{U}_1$ (constructing a global retraction seems difficult).

Both spaces have homotopy type of CW -complexes and to prove homotopy equivalence it is enough to prove that any continuous map $f : S^k \rightarrow \overline{\mathcal{U}}_1$ (where S^k is a sphere) may be homotoped to a map $S^k \rightarrow \mathcal{U}_1$. Assume that there is a tunnel $T = (y, z, L)$ such that $(T / \sim) \in f(S^k) \cap (\overline{\mathcal{U}}_1 - \mathcal{U}_1)$. We have $L > 0$. For any $\varepsilon > 0$, let $B_\varepsilon(y)$ be the open ball of radius ε in Y around y , let $B_\varepsilon(z)$ be the affine interval $(z - \sqrt{-1}\varepsilon, z + \sqrt{-1}\varepsilon)$. For ε small enough, there is a unique continuous function $L_\varepsilon : B_\varepsilon(y) \times B_\varepsilon(z) \rightarrow \mathbb{R}$ such that $L_\varepsilon(y, z) = L$ and for all $(y', z') \in B_\varepsilon(y) \times B_\varepsilon(z)$, $(y', z', L_\varepsilon(y', z'))$ represents a point in $\overline{\mathcal{U}}_1 - \mathcal{U}_1$. The “half-ball”

$$H_\varepsilon := \{(y', z', L') \in B_\varepsilon(y) \times B_\varepsilon(z) \times \mathbb{R} \mid 0 \leq L' \leq L_\varepsilon(y', z')\}$$

is a neighbourhood of $T / \sim$ in $\overline{\mathcal{U}}_1$. Working inside this neighbourhood, one may homotope f to f' such that $f'(S^k) \cap (\overline{\mathcal{U}}_1 - \mathcal{U}_1) \subseteq f(S^k) \cap (\overline{\mathcal{U}}_1 - \mathcal{U}_1) - \{T / \sim\}$. Compacity of $f(S^k) \cap (\overline{\mathcal{U}}_1 - \mathcal{U}_1)$ guarantees that one can iterate this process a finite number of times to get rid of all $f(S^k) \cap (\overline{\mathcal{U}}_1 - \mathcal{U}_1)$. $\square$

Proposition 9.6. *The nerve of $(\widehat{\mathcal{U}}_b)_{b \in B}$ is $\text{Flag}(B, S)$.*

Proof. Let $b, b' \in B$ such that $\widehat{\mathcal{U}}_b \cap \widehat{\mathcal{U}}_{b'} \neq \emptyset$. Let $T = (y, x, L)$ and $T' = (y', x', L')$ be semitunnels, representing points z and z' in $\widehat{\mathcal{U}}_1$, such that $bz = b'z'$. The image of z (resp. z') in $W \setminus V^{\text{reg}}$ is $(x + L, y)$ (resp. $(x' + L', y')$). Thus $x + L = x' + L'$ and $y = y'$. Up to permuting b and b' , we may assume that $L \geq L'$. Since $x' = x + L - L'$ is in U_y , $T'' := (y, x, L - L')$ is a tunnel, representing a simple element b'' . The tunnel T is a concatenation of T'' and T' . This implies that $z = b''z'$ and $b'z' = bz = bb''z'$. By faithfulness of the B -action on the orbit of z , we conclude that $bb'' = b'$.

We have proved that the 1-skeletons of the nerve and of $\text{Flag}(B, S)$ coincide. To conclude, it remains to check that the nerve is a flag complex. Let $C \subseteq B$ be such that for all $b, b' \in C$, either $b^{-1}b'$ or $b'^{-1}b$ is simple. We have to prove that $\bigcup_{b \in C} \widehat{\mathcal{U}}_b \neq \emptyset$.

Let $c_0, \dots, c_m$ be the elements of C , numbered according to the total ordering on C induced by $\preceq$:

$$c_0 \prec c_1 \prec c_2 \prec \dots \prec c_m \preceq c_0 \delta.$$

Up to left-dividing each term by c_0 , we may assume that $c_0 = 1$. Let $y \in Y_C$. We may find $x, L_1, \dots, L_m$ such that (y, x, L_i) represents c_i for all i . The point of the universal cover represented by (y, x, L_m) belongs to $\bigcup_{b \in C} \widehat{\mathcal{U}}_b$. $\square$

Proposition 9.7. *Let C be a subset of B such that $\bigcap_{b \in C} \widehat{\mathcal{U}}_b \neq \emptyset$. Then $\bigcap_{b \in C} \widehat{\mathcal{U}}_b$ is contractible.*

Proof. As in the previous proof, we write $C = \{c_0, c_1, \dots, c_m\}$ with

$$c_0 \prec c_1 \prec c_2 \prec \dots \prec c_m \preceq c_0 \delta$$

and assume without loss of generality that $c_0 = 1$.

The case $m = 0$ is Lemma 9.5.

Assume that $m \geq 1$. Let $T = (y, z, L)$. The point represented by T lies in $\bigcap_{b \in C} \widehat{\mathcal{U}}_b$ if and only if there exists $L_1, \dots, L_m$ with $0 < L_1 < \dots < L_m < L$ such that, for all i , $T_i := (y, z, L_i)$ is a tunnel representing c_i . Given $y \in Y$, it is possible to find such $L_1, \dots, L_m$ if and only if $y \in Y_C$. This justifies:

$$\bigcap_{b \in C} \widehat{\mathcal{U}}_{b,y} \neq \emptyset \Leftrightarrow y \in Y_C.$$

For a given $y \in Y_C$, let us study the intersection $\bigcap_{b \in C} \widehat{\mathcal{U}}_{b,y}$. Let $(x_1, \dots, x_k)$ be the ordered support of $LL(y)$. Let $A_0, \dots, A_m$ be the associated partition of $LL(y)$, defined in Proposition 8.4. Let

$$\begin{aligned} R_+(y, C) &:= \max\{\text{re}(z) \mid z \in A_m\}, \\ R_-(y, C) &:= \min\{\text{re}(z) \mid z \in A_1\}, \\ I_+(y, C) &:= \min\{\text{im}(z) \mid z \in A_1 \cup \dots \cup A_m\}, \end{aligned}$$


and

$$I_-(y, C) = \sup\{\text{im}(z) \mid z \in A_0 \text{ and } R_- \leq \text{re}(z) \leq R_+\}.$$

We have $I_-(y, C) < I_+$. It may happen that $I_-(y, C) = -\infty$.

We illustrate this on a picture. In the example, the support of $LL(y)$ is $x_1, \dots, x_6$ and $C = \{1 \prec s_2 \prec s_2 s_4\}$, where $(s_1, \dots, s_6) = \text{lbl}(y)$. The support of A_1 is x_2 and the support of A_2 is x_4 . The remaining points are in A_0 . The lines $\text{im}(z) = I_-(y, C)$ and $\text{im}(z) = I_+(y, C)$ are represented by full lines. A semitunnel (y, z, L) representing

a point in $\bigcap_{b \in C} \widehat{\mathcal{U}}_{b,y}$ must cross the intervals I_2 and I_4 represented by dotted lines. One must have $\operatorname{re}(z) < R_-(y, C)$ and $I_-(y, C) < \operatorname{im}(z) < I_+(y, C)$; the final point $z + L$ must satisfy $\operatorname{re}(z + L) > R_+(y, C)$. This final point may be any complex number z' in the rectangle $\operatorname{re}(z') > R_+(y, C)$ and $I_-(y, C) < \operatorname{im}(z') < I_+(y, C)$, except the points on the closed horizontal half-line to the right of x_6 (indicated by a dashed line), which cannot be reached.



Generalising the example, one shows that $\bigcap_{b \in C} \widehat{\mathcal{U}}_b$ may be identified with

$$\bigcup_{y \in Y_C} E(y, C),$$

where $E(y, C)$ is the open rectangle of $\mathbb{C}$ defined by

$$\operatorname{re}(z) > R_+(y, C), \quad I_-(y, C) < \operatorname{im}(z) < I_+(y, C),$$

from which have been removed the possible points of A_0 and the horizontal half-lines to their rights.

Let $\overline{E}(y, C)$ be the rectangle

$$\operatorname{re}(z) \geq R_+(y, C), \quad I_-(y, C) < \operatorname{im}(z) < I_+(y, C)$$

from which have been removed the possible points of A_0 and the horizontal half-lines to their rights.

A homotopy argument similar to the one used in the proof of Lemma 9.5 (or possibly a nicer argument from semialgebraic geometry) shows that $\bigcup_{y \in Y_C} E(y, C)$ and $\bigcup_{y \in Y_C} \overline{E}(y, C)$ are homotopy equivalent. The latter may be retracted to the union of open intervals $\bigcup_{y \in Y_C} (I_-(y, C), I_+(y, C))$ (on each rectangle, the retraction is

$$\overline{E}(y, C) \times [0, 1] \rightarrow \overline{E}(y, C), (z, t) \mapsto R_+(y, C) + t(\operatorname{re}(z) - R_+(y, C)) + \sqrt{-1} \operatorname{im}(z).$$

The union $\bigcup_{y \in Y_C} (I_-(y, C), I_+(y, C))$ may be retracted to $\bigcup_{y \in Y_C} [I_0(y, C), I_+(y, C))$, where $I_0(y, C) := \max(\frac{I_-(y, C) + I_+(y, C)}{2}, I_+(y, C) - 1)$. The latter space is a fibre bundle over Y_C , with fibers intervals. Since the base space Y_C is contractible (Proposition 8.3), this fibre bundle is contractible. So is $\bigcap_{b \in C} \widehat{\mathcal{U}}_b$. $\square$

10. GENERALISED NON-CROSSING PARTITIONS

Here again, W is an irreducible well-generated complex 2-reflection group.

When W is of type A_{n-1} , the lattice $(S, \preceq)$ is isomorphic to the lattice of non-crossing partitions of a regular n -gon ([6], [10]). Following [30], [3] and [5], we call *generalised lattice of non-crossing partitions of type W* the lattice

$$(S, \preceq),$$

and *Catalan number of type W* the number

$$\text{Cat}(W) := \prod_{i=1}^n \frac{d_i + h}{d_i}.$$

The operation sending $s \preccurlyeq t$ to $s^{-1}t$ is an analog of Kreweras complement operation. The map $s \mapsto s^{-1}\delta$ is an anti-automorphism of the lattice.

In the Coxeter case, Chapoton discovered a general formula for the number $Z_W(N)$ of weak chains $s_1 \preccurlyeq s_2 \preccurlyeq \cdots \preccurlyeq s_N$ of length N in $(S, \preccurlyeq)$ (see [15]). This formula continues to hold, though we are only able to prove this case-by-case (see [1] and [15] for the Coxeter types; the $G(e, e, n)$ case was done in [5]; the remaining types are done by computer).

Proposition 10.1. *We have, for all N ,*

$$Z_W(N) = \prod_{i=1}^n \frac{d_i + Nh}{d_i}.$$

Corollary 10.2. *We have $|S| = \text{Cat}(W)$.*

Another interesting numerical invariant is the *Poincaré polynomial*

$$\text{Poin}(S) := \sum_{s \in S} t^{l(s)}.$$

The table below summarises the numerical data for the exceptional types (real and non-real). The coefficient of t in the Poincaré polynomial is the cardinal of R_c . One observes that

$$R = R_c \Leftrightarrow W \text{ is real.}$$

In the Weyl group case, $\text{Poin}(S)$ may be interpreted as the Poincaré polynomial of the cohomology of a toric variety related to cluster algebras ([15]).

W	degrees	$ R $	$\text{Cat}(W)$	$\text{Poin}(S)$	$ \text{Red}_T(c) $
$G_{23} (H_3)$	2, 6, 10	15	32	$1 + 15t + 15t^2 + t^3$	50
G_{24}	4, 6, 14	21	30	$1 + 14t + 14t^2 + t^3$	49
G_{27}	6, 12, 30	45	42	$1 + 20t + 20t^2 + t^3$	75
$G_{28} (F_4)$	2, 6, 8, 12	24	105	$1 + 24t + 55t^2 + 24t^3 + t^4$	432
G_{29}	4, 8, 12, 20	40	112	$1 + 25t + 60t^2 + 25t^3 + t^4$	500
$G_{30} (H_4)$	2, 12, 20, 30	60	280	$1 + 60t + 158t^2 + 60t^3 + t^4$	1350
G_{33}	4, 6, 10, 12, 18	45	308	$1 + 30t + 123t^2 + 123t^3 + 30t^4 + t^5$	4374
G_{34}	6, 12, 18, 24, 30, 42	126	1584	$1 + 56t + 385t^2 + 700t^3 + 385t^4 + 56t^5 + t^6$	100842
$G_{35} (E_6)$	2, 5, 6, 8, 9, 12	36	833	$1 + 36t + 204t^2 + 351t^3 + 204t^4 + 36t^5 + t^6$	41472
$G_{36} (E_7)$	2, 6, 8, 10, 12, 14, 18	63	4160	$1 + 63t + 546t^2 + 1470t^3 + 1470t^4 + 546t^5 + 63t^6 + t^7$	1062882
$G_{37} (E_8)$	2, 8, 12, 14, 18, 20, 24, 30	120	25080	$1 + 120t + 1540t^2 + 6120t^3 + 9518t^4 + 6120t^5 + 1540t^6 + 120t^7 + t^8$	37968750

11. ON BADLY-GENERATED GROUPS

Our theory works with well-generated groups, but we conjecture that the dual braid monoid is a particular case of a more general construction, giving good monoids for braid groups of badly-generated groups. If $d = d_{i_0}$ is a regular degree of an arbitrary irreducible complex reflection group, let L be a generic line of direction X_{i_0} in $W \setminus V$ (as in [2, section 3]). It intersects the discriminant variety $\mathcal{H}$ in $(N + N^*)/d$ points. Let C be an Euclidean circle in L surrounding this intersection. It represents a “Coxeter element” in the free group $\pi_1(L - L \cap \mathcal{H})$, as in [4], and we have a natural “dual braid monoid” for the free group, generated by non-crossing loops drawn inside C . We conjecture that the direct image of this monoid in $B(W)$ is a *quasi-Garside monoid*, i.e., it satisfies all axioms of Garside monoids except the finiteness of the set of simples (the finiteness should occur only if W is well-generated and $d = h$ – it is not difficult to see that, in this case, this monoid coincides with the one studied in the present article). The main difficulty is that this prevents the use of a counting argument such as the one used in this article.

REFERENCES

- [1] Ch. Athanasiadis, V. Reiner, *Non-crossing partitions for the group D_n* , preprint.
- [2] D. Bessis, *Zariski theorems and diagrams for braid groups*, Invent. Math. **145** (2001), 487–507.
- [3] D. Bessis, *The dual braid monoid*, Ann. Sci. École Norm. Sup. **36** (2003), 647–683.
- [4] D. Bessis, *A dual braid monoid for the free group*, arXiv:math.GR/0401324.
- [5] D. Bessis, R. Corran, *Non-crossing partitions of type (e, e, r)* , arXiv:math.GR/0403400.
- [6] D. Bessis, F. Digne, J. Michel, *Springer theory in braid groups and the Birman-Ko-Lee monoid*, Pacific J. Math. **205** (2002), 287–310.
- [7] D. Bessis, J. Michel, *Explicit presentations for exceptional braid groups*, to appear in Exp. Math.
- [8] M. Bestvina, *Non-positively curved aspects of Artin groups of finite type*, Geometry and Topology **3** (1999), 269–302.
- [9] J. Birman, K. H. Ko, S. J. Lee, *A new approach to the word and conjugacy problem in the braid groups*, Adv. Math. **139** (1998), no. 2, 322–353.
- [10] T. Brady, *A partial order on the symmetric group and new $K(\pi, 1)$ ’s for the braid groups*, Adv. Math. **161** (2001), 20–40.
- [11] T. Brady, C. Watt, *$K(\pi, 1)$ ’s for Artin groups of finite type*, Geom. Dedicata **94** (2002), 225–250.
- [12] E. Brieskorn, *Die Fundamentalgruppe des Raumes der regulären Orbits einer endlichen komplexen Spiegelungsgruppe*, Invent. Math. **12** (1971), 57–61.
- [13] E. Brieskorn, *Sur les groupe de tresses*, Séminaire Bourbaki 1971/72, Lectures Notes in Math. **317**, Springer-Verlag, 1973, 21–44.
- [14] M. Broué, G. Malle, R. Rouquier, *Complex reflection groups, braid groups, Hecke algebras*, J. reine angew. Math. **500** (1998), 127–190.
- [15] F. Chapoton, *Enumerative properties of generalized associahedra*, Sémin. Lothar. Combin. **51** (2004), Art. B51b, 16 pp. (electronic).
- [16] R. Charney, J. Meier, K. Whittlesey, *Bestvina’s normal form complex and the homology of Garside groups*, Geom. Dedicata **105** (2004), 171–188.
- [17] W. Couwenberg, G. Heckman, E. Looijenga, *Geometric structures on the complement of a projective arrangement*, arXiv:math.AG/0311404.
- [18] P. Dehornoy, *Groupes de Garside*, Ann. Sci. École Norm. Sup. (4) **35** (2002) 267–306.
- [19] P. Dehornoy, L. Paris, *Gaussian groups and Garside groups, two generalizations of Artin groups*, Proc. of London Math. Soc. **79** (1999), 569–604.
- [20] P. Deligne, *Les immeubles des groupes de tresses généralisés*, Invent. Math. **17** (1972), 273–302.
- [21] P. Deligne, letter to E. Looijenga, 9/3/1974.
- [22] E. Fadell, L. Neuwirth, *Configuration spaces*, Math. Scand. **10** (1962), 111–118.

- [23] A. Hatcher, *Algebraic topology*, Cambridge University Press, 2002.
- [24] G. Lehrer, J. Michel, *Invariant theory and eigenspaces for unitary reflection groups*, C. R. Math. Acad. Sci. Paris **336** (2003), no. 10, 795–800.
- [25] E. Looijenga, *The complement of the bifurcation variety of a simple singularity*, Invent. Math. **23** (1974), 105–116.
- [26] T. Nakamura, *A note of the $K(\pi, 1)$ -property of the orbit space of the unitary reflection group $G(m, l, n)$* , Sci. Papers College of Arts and Sciences, Univ. Tokyo **33** (1983), 1–6.
- [27] P. Orlik, L. Solomon, *Unitary reflection groups and cohomology*, Invent. Math. **59** (1980), 77–94.
- [28] P. Orlik, L. Solomon, *Discriminants in the invariant theory of reflection groups*, Nagoya Math. J. **109** (1988), 23–45.
- [29] P. Orlik, H. Terao, *Arrangements of hyperplanes*, Grundlehren der mathematischen Wissenschaften **300**, Springer-Verlag, 1992.
- [30] V. Reiner, *Non-crossing partitions for classical reflection groups*, Discrete Math. **177** (1997), 195–222.
- [31] K. Saito, *On a linear structure of the quotient variety by a finite reflection group*, Publ. Res. Inst. Math. Sci. Kyoto Univ. **29** (1993), 535–579.
- [32] K. Saito, *Uniformization of the orbifold of a finite reflection group*, RIMS preprint, may 2003.
- [33] K. Saito, *The polyhedra dual to the Weyl chamber decomposition: a précis*, RIMS preprint, september 2004.
- [34] T. A. Springer, *Regular elements of finite reflection groups*, Invent. Math. **25** (1974), 159–198.
- [35] G. C. Shephard, J. A. Todd, *Finite unitary reflection groups*, Canad. J. Math. **6** (1954), 274–304.

DMA, ÉCOLE NORMALE SUPÉRIEURE, 45 RUE D’ULM, 75230 PARIS CEDEX 05, FRANCE
E-mail address: david dot bessis at ens dot fr